

# The Tree of Sex consortium: a global initiative for studying the evolution of reproduction in eukaryotes

Daniel Jeffries<sup>1</sup>, Chiara Benvenuto<sup>2</sup>, Astrid Böhne<sup>3</sup>, Christelle Fraïsse<sup>4</sup>, Sònia Garcia<sup>5</sup>, Paul Jay<sup>6</sup>, Lukáš Kratochvíl<sup>7</sup>, Caitlin E. McDonough-Goldstein<sup>8</sup>, Aurora Ruiz-Herrera<sup>9,10</sup>, Cibeles G. Sotero-Caio<sup>11</sup>, Nicole Valenzuela<sup>12</sup>, Melissa A. Wilson<sup>13,14,15</sup>, Tree of Sex Consortium, Kamil S. Jaron<sup>11</sup>

<sup>1</sup>Division of Evolutionary Ecology, Institute of Ecology and Evolution, University of Bern, Bern, Switzerland

<sup>2</sup>Environmental Research and Innovation Centre (ERIC), School of Science, Engineering and Environment, University of Salford, Salford, United Kingdom

<sup>3</sup>Centre for Molecular Biodiversity Research, Leibniz Institute for the Analysis of Biodiversity Change LIB, Museum Koenig Bonn, Bonn, Germany

<sup>4</sup>CNRS, Univ. Lille, UMR 8198 – Evo-Eco-Paleo, Lille, France

<sup>5</sup>Institut Botànic de Barcelona, IBB (CSIC-CMNCB), Barcelona, Spain

<sup>6</sup>Center for GeoGenetics, University of Copenhagen, Copenhagen, Denmark

<sup>7</sup>Department of Ecology, Faculty of Science, Charles University, Prague, Czech Republic

<sup>8</sup>Department of Integrative Biology, University of Wisconsin-Madison, Wisconsin, United States of America

<sup>9</sup>Departament de Biologia Cel·lular, Fisiologia i Immunologia, Universitat Autònoma de Barcelona, Cerdanyola del Vallès, Spain

<sup>10</sup>Genome Integrity and Instability Group, Institut de Biotecnologia i Biomedicina, Universitat Autònoma de Barcelona, Cerdanyola del Vallès, Spain

<sup>11</sup>Tree of Life, Wellcome Sanger Institute, Wellcome Genome Campus, Cambridge, United Kingdom

<sup>12</sup>Department of Ecology, Evolution, and Organismal Biology, Iowa State University, Ames, IA, United States

<sup>13</sup>School of Life Sciences, Arizona State University, Tempe, AZ, United States

<sup>14</sup>Center for Evolution and Medicine, Arizona State University, Tempe, AZ, United States

<sup>15</sup>Center for Mechanisms of Evolution, Biodesign Institute, Tempe, AZ, United States

Handling editor: Richard Meisel, Guest editor: Sylvain Glemin

Corresponding authors: Daniel Jeffries, Division of Evolutionary Ecology, Institute of Ecology and Evolution, University of Bern, Baltzerstrasse 6, 3012 Bern, Switzerland. Email: [djeffries86@gmail.com](mailto:djeffries86@gmail.com); Kamil S. Jaron, Tree of Life, Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridge, CB10 1SA, UK. Email: [kj11@sanger.ac.uk](mailto:kj11@sanger.ac.uk)

## Abstract

Reproduction is a fundamental aspect of life that affects all levels of biology, from genomes and development to population dynamics and diversification. The first Tree of Sex database synthesized a vast diversity of reproductive strategies and their intriguing distribution throughout eukaryotes. A decade on, we are reviving this initiative and greatly expanding its scope to provide the most comprehensive integration of knowledge on eukaryotic reproduction to date. In this perspective, we first identify important gaps in our current knowledge of reproductive strategies across eukaryotes. We then highlight a selection of questions that will benefit most from this new Tree of Sex project, including those related to the evolution of sex, modes of sex determination, sex chromosomes, and the consequences of various reproductive strategies. Finally, we outline our vision for the new Tree of Sex database and the consortium that will create it ([treeofsex.org](http://treeofsex.org)). The new database will cover all Eukaryota and include a wide selection of biological traits. It will also incorporate genomic data types that were scarce or non-existent at the time of the first Tree of Sex initiative. The new database will be publicly accessible, stable, and self-sustaining, thus greatly improving the accessibility of reproductive knowledge to researchers across disciplines for years to come. Lastly, the consortium will persist after the database is created to serve as a collaborative framework for research, prioritizing ethical standards in the collection, use, and sharing of reproductive data. The new Tree of Sex consortium is open, and we encourage all who are interested in this topic to join us.

**Keywords:** biodiversity, asexual reproduction, sexual reproduction, sex chromosomes, sex determination, database, ontology

## Introduction

Reproduction is a defining characteristic of life. Whether sexual, clonal, or one of a myriad of strategies in between, every extant species has some successful method for propagation. Importantly, an organism's **reproductive strategy** (see the glossary for all terms in bold) can influence all biological levels of organization, from genomes, cellular processes, development,

physiology, through populations, ecology, behaviour, genetic diversity and the propensity for adaptation, diversification, and extinction. Describing the diversity of reproductive strategies among organisms and investigating how and why they evolve is thus essential when trying to understand the natural world.

Given the numerous biological factors associated with reproductive strategies, their characterization in diverse taxa

Received August 10, 2024; revised April 18, 2025; accepted May 1, 2025

© The Author(s) 2025. Published by Oxford University Press on behalf of the European Society of Evolutionary Biology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

**Table 1.** Example of existing databases containing data relevant to reproduction. The species numbers were derived from publications or directly by counting unique species names within available databases. Except for Tree of Sex v1, the databases are listed in alphabetical order.

Database name	Scope	Taxonomic scope	Species number	Reference
Tree Of Sex v1	Sexual systems, sex chromosomes	Vertebrates, invertebrates, plants	37,496	(Tree of Sex Consortium, 2014)
Animal Chromosome Count, ACC	Chromosome counts	Animals	14,524	(Román-Palacios et al., 2021)
AmphibianKaryo	Sex chromosomes, chromosome counts	Amphibians	2,124	(Perkins et al., 2019)
AndrodioecyAnimal	Androdioecy	Animal	36	(Weeks et al., 2006)
Animal_rDNA	Chromosome counts, karyotypes, genome size	Animals	2,800	(Sochorová et al., 2018)
AnimalGenomeSize	Genome size (C-values)	Animals	6,534	(Gregory, 2024)
ASER	Sex reversal	Animals	18	(Y. Li et al., 2021)
B-chrom	B chromosomes	Animals, plants, fungi	2,951	(D'Ambrosio et al., 2017; Gutiérrez et al., 2023)
CCDB	Chromosome counts	Plants	77,958	(Rice et al., 2015)
ChromNematoda	Chromosome counts, reproductive strategy	Nematodes	257	(Blaxter et al., 2024a)
ColeopteraKaryo	Sex chromosomes, chromosome counts	Beetles	4,960	(Blackmon & Demuth, 2015a)
DipteraKaryo	Sex chromosomes, chromosome counts	Diptera	3,443	(Morelli et al., 2022)
DrosophilaKaryo	Sex chromosomes, chromosome counts	Drosophila	1,247	(Morelli et al., 2022)
FishBase	Reproduction mode, life-history traits	Fish	35,400	(Froese & Pauly, 2024)
FISHKARYOME	Sex chromosomes, chromosome counts	Fish	1,285	(Nagpure et al., 2016)
FishSexChrom	Sex chromosomes (direct comparison with ToS v.1)	Fish	440	(Sember et al., 2021)
HarvestmenCyto	Chromosome counts	Harvestmen	100	(Tsurusaki et al., 2022)
HerpSexDet	Sex determination, sex reversal	Amphibians, reptiles	891	(Nemesházi & Bókony, 2023)
HZmatingSystem	Mating systems	Plants (hybrid zones)	245	(Pickup et al., 2019, 2020)
Insect_Egg_Evolution	Egg size and shape	Insects	6,700	(Church et al., 2019)
InSexBase	Sex chromosomes, sex-biased genes	Insects	49	(Chen et al., 2021)
KaryoBlatto	Chromosome counts	Termites, cockroaches	229	(Jankásek et al., 2021)
Mammal_SameSexSexual-Behavior	Observed same-sex sexual behavior	Mammals	322	(Gómez et al., 2023)
MammalKaryo	Sex chromosomes, chromosome counts	Mammals	1,440	(Blackmon et al., 2019)
NESCent	Mating systems	Plants, life-history traits	154	(Goodwillie et al., 2005b, 2010)
Phaeodev	Genome sizes, ploidy	Brown algae	67	(The Phaeoexplorer project, 2024)
Phycosm	Genome sizes	Algae, heterokonts	154	(Grigoriev et al., 2021)
Plant_DNA_C-values	Genome sizes (c-values), chromosome counts	Plants	12,273	(Leitch et al., 2019; Pelli- cer & Leitch, 2020)
Plant_rDNA	Chromosome counts, karyotypes, genome size	Plants	2,770	(Rodríguez-González et al., 2023)
PolyneopteraKaryo	Sex chromosomes, chromosome counts	Polyneoptera	823	(Sylvester et al., 2020)
PseudoscorpionCyto	Chromosome counts	Pseudoscorpions	65	(Štáhlavský, 2022)
ReprodTraitsAlgae	Sexual systems, life-history traits	Brown algae	91	(Heesch et al., 2021)
ROSIE	Sex determination	Testudines	125	(Krueger & Janzen, 2022)
SAGD	Sex-biased genes	Animals	21	(Shi et al., 2019)

Table 1. Continued

Database name	Scope	Taxonomic scope	Species number	Reference
ScorpionCyto	Chromosome counts	Scorpions	264	(Schneider et al., 2024)
Sex-chrom	Sex chromosomes, ploidy, chromosome counts	Land plants, green algae	229	(Baránková et al., 2020; Garcia et al., 2023)
SexChromAlgae	Sex chromosomes	Brown algae	10	(Barrera-Redondo et al., 2024)
SexSystemCrustacea	Sexual systems	Crustaceans	334	(Benvenuto & Weeks, 2020)
SexSystemFish	Sexual systems	Fish	4,614	(Pla et al., 2022)
SpermTree	Sexual systems, sperm morphology	Animals	5,675	(Fitzpatrick et al., 2022)
SpiderCyto	Chromosome counts	Spiders	933	(Araujo et al., 2024)

requires multiple approaches and expertise from a broad range of disciplines, including genetics/genomics, cell biology, physiology, developmental biology, ecology, behavioural ecology, and evolutionary biology. Understanding the drivers and consequences of these strategies throughout nature requires subsequent in-depth analyses against a reliable phylogeny. However, before this is possible, information from disparate methodological and taxonomic disciplines must be integrated.

Roughly a decade ago, a group of scientists created the Tree of Sex database (Tree of Sex Consortium, 2014), collected data for over 37,000 species, across many eukaryotic lineages, and used it to gain several foundational insights into the distribution and evolution of reproductive strategies. These included the previously unrecognized lability and patterning of sex determination mechanisms across eukaryotes (Bachtrog et al., 2014), and well-supported estimates of variation in the rates of transitions among different reproductive strategies (Blackmon & Demuth, 2014; Blackmon et al., 2017; Pennell et al., 2018). Since then, extensive research and methodological advancements, including the continuing genomic revolution, have deepened our insights into reproductive biology. To capitalize on these advances, we have revived the Tree of Sex consortium. We are now working towards a new database that not only updates the information compiled by the first initiative, but also largely expands its scope to include all clades of eukaryotes for which data exists, all reproductive strategies identified to date, and new data types, including genomic data. Crucially, we will integrate this knowledge via a purpose-built ontology for reproductive biology, allowing us to connect information from disparate fields and from studies that are decades apart. The new Tree of Sex consortium will also serve as a framework for collaborative research to address fundamental questions about the evolution of reproductive strategies.

To set the stage for the new Tree of Sex project, here we summarize existing efforts to describe the diversity of eukaryotic reproductive strategies and highlight the current gaps in our knowledge and their importance. We then describe some of the topics and unanswered questions that we believe will benefit most from large comparative approaches that the Tree of Sex database will enable. Lastly, we describe our plans for the Tree of Sex database and the consortium, emphasizing the need for proactive engagement of a diversity of people, understandings, and perspectives so that we can build a database that encompasses the complexity and nuances of reproduction and its contexts throughout nature.

## Characterizing the diversity of reproductive strategies across eukaryotes

### Existing data on eukaryotic reproduction

Taxonomically speaking, the first Tree of Sex database (Tree of Sex Consortium, 2014) remains the broadest integrated collection of information on reproduction in eukaryotes compiled to date. In the decade since, several additional databases (Table 1) have been assembled, focusing on reproduction-related traits for distinct groups of organisms, including (but not limited to) fishes (Pla et al., 2022; Sember et al., 2021), amphibians and reptiles (Nemesházi & Bókony, 2023), insects (Chen et al., 2021), and plants (Baránková et al., 2020; Garcia et al., 2023). In addition, several other databases contain information that is relevant but not necessarily specific to reproduction, including karyotypes (Blackmon & Demuth, 2015a; Blaxter et al., 2024a; D'Ambrosio et al., 2017; Garcia et al., 2012; Jankásek et al., 2021; Román-Palacios et al., 2021; Sochorová et al., 2018) and genome sizes (Gregory, 2024; Pellicer & Leitch, 2020). Because reproductive strategies depend on and shape genomic characteristics (e.g., divergence, sex chromosomes, karyotype), these data types can greatly improve our understanding of the reproductive strategy of a given species.

Together, the above databases contain information for tens of thousands of species, including the presence or absence of sexual reproduction, mating systems, sex allocation, sex determination, karyotype, sex chromosome differentiation, ecological and behavioural phenotypes and many other traits that can inform the study of reproduction. However, these datasets are currently difficult to use in combination, as they lack standardized terms. Moreover, the lack of a common phylogeny hampers evolutionary analyses outside of the taxonomic bounds of each individual database.

### Gaps in our knowledge of reproduction in eukaryotes

Despite extensive efforts, existing databases capture only a portion of the existing literature, which itself does not encompass the full diversity of reproductive strategies in nature. To identify taxonomic gaps in our knowledge, we conducted a literature search based on a non-exhaustive set of 36 keywords related to reproductive strategies (see legend of Figure 1). The search hits reveal a substantial taxonomic bias in the literature on reproductive strategies (Figure 1A). Unsurprisingly, species with relevance to medicine, agriculture, and basic

research (such as humans, mice, fruit flies, and the roundworm *Caenorhabditis elegans*) have received the most research interest. Indeed, over half of all papers studying chordates focus on humans or mice. Consequently, the spread of research across individual phyla does not reflect their species richness (Figure 1C).

One additional factor likely contributing to these taxonomic biases is the difficulty of characterizing reproductive strategies in non-model organisms. However, technological advancements are mitigating many such issues. Genomic techniques can help characterize the ploidy, karyotype, inheritance, sex chromosomes, sex-specific gene expression patterns, population dynamics, and other reproduction-relevant traits, in many cases without any prior knowledge. While it is still necessary to integrate such data with ecological, phenotypic, quantitative genetics, and cytogenetic approaches in order to gain a holistic picture of reproduction in a given species, genomics can provide a valuable foundation on which to build our understanding of reproduction, even in hard to study organisms (Deakin et al., 2019; Stöck et al., 2021a; Zaidem et al., 2019).

In addition to biases from methodological limitations or applied practices, it is well known that the attention received by a given study system in biology is also affected by cultural and geographical biases. Owing to differences in resources and funding access, the most biodiverse regions of the planet have received the least research focus (Linck & Cadena, 2024; Titley et al., 2017). This also affects the diversity of researchers studying a given species as well as phrasing of hypotheses (Ahnesjö et al., 2020). Despite the ever-increasing yearly publication rates across science (Bornmann et al., 2021), the taxonomic biases in the distribution of publications on reproductive strategies per phylum have remained stable for over 70 years (Figure 1A). The available genomes show a different skew (Figure 1B) to that of the published literature, however, this has again changed little through time. Together, the stability of these trends highlights that closing the taxonomic gaps in our knowledge will require proactivity in the choice of future study systems and a serious push for inclusive research strategies that provide opportunities to researchers in low- and middle-income countries.


When defining a strategy to mitigate problems caused by existing biases in study systems, one important factor to consider is the reproductive diversity of the studied taxa. For example, the large majority of studied viviparous mammals possess an XX/XY sex determination system with Sry as a sex determining gene, which has remained conserved for hundreds of millions of years (Hughes & Page, 2015). As such, it is less likely that exploring additional viviparous mammal systems will significantly broaden our knowledge of diversity and frequency of sex determination systems in eukaryotes. In contrast, based on the species studied so far, animals such as amphibians (Jeffries et al., 2018; Ma & Veltsos, 2021), reptiles (Gamble et al., 2015; Holleley et al., 2015; Krueger & Janzen, 2022; Pinto et al., 2022, 2023), fish (El Taher et al., 2021; Jeffries et al., 2018; Kitano et al., 2024), crustaceans (J. Li, 2022; Ye et al., 2023), insects (Blackmon et al., 2017; Bracewell et al., 2024; Vicoso & Bachtrog, 2015), arachnids (Araujo et al., 2012; Cordellier et al., 2020; Kořínková & Král, 2013), and potentially molluscs (Breton et al., 2018; Yusa, 2007) seem to display a large diversity of sex determination mechanisms, and yet they have received comparatively little attention in this regard. Similarly, haploid sexual systems

are severely understudied, despite being found in many taxa including algae and bryophytes (Charlesworth, 2022; Coelho et al., 2018) and being important for testing predictions about sex chromosome evolution. Finally, there are many more fundamental gaps in our knowledge of reproduction of unicellular eukaryotes. For example, in microsporidia (important spore-forming unicellular fungi that parasitize all major groups of animals) all we know is that some form of sexual reproduction likely occurs, based on the presence of an intact multi-gene cassette involved in mating-type determination in fungi (Lee et al., 2008), and putatively imaged gametes (reviewed in Khalaf et al., 2024). These taxa, which have traditionally been overlooked and show signs of high variability in reproductive strategies, likely hold the most potential for broadening our knowledge of reproductive strategy diversity and understanding the relative frequencies of the many systems that exist in nature.

We note, however, that we are not petitioning against the further study of already-well-studied clades. Indeed, the ideal study system is defined by the question at hand, and the in-depth study of model systems, and those that are exceptions to clade norms can be extremely insightful for specific evolutionary processes, as has been the case with the few exceptional mammalian species which do not share the usual mammalian XX/XY sex determination system (e.g., Fredga, 1988; Hughes et al. 2024 ; Mulugeta et al., 2016; Ruiz-Herrera & Waters, 2022; Saunders & Veyrunes, 2021).

Aside from taxonomic gaps, the types of data available for a given study system are also sporadic, with many lacking cytological (e.g., chromosome morphologies, centromere positions) or genomic data. The combination of high-throughput genomics with cytogenetic analyses (as in Deakin et al., 2019; Liehr, 2021) will undoubtedly prove to be a powerful approach to studying patterns of chromosomal rearrangements, such as inversions, translocations, fusions, or fissions, which can all impact reproductive processes including recombination in meiosis or gametogenesis, e.g., resulting in complete versus incomplete genome complements in gametes (Baránková et al., 2020; Deakin et al., 2019; Gil-Fernández et al., 2020; Vara et al., 2021), as well as speciation and hybridization. Such approaches will also be instrumental in describing phenomena such as programmed DNA elimination, which leads to differences in (sex) chromosome numbers or synteny between germline and somatic genomes. While **programmed DNA elimination** has so far been observed in multiple eukaryotic lineages (Hodson & Ross, 2021; Smith et al., 2021; Wang & Davis, 2014), its frequency is still not well known.

Advancements in chromosome conformation capture (“3C”) methods, such as Hi-C (Jerkovic & Cavalli, 2021; Rao et al., 2014), are also opening new avenues for studying genome evolution across various timescales and cell types (Álvarez-González et al., 2022; Bista et al., 2024; Hoencamp et al., 2021; Vara et al., 2019). This technology has already provided insights into the chromatin structure of sex chromosomes in both somatic and germ cells, albeit in a limited number of organisms (Bista et al., 2024; Chen et al., 2020; Hoencamp et al., 2021; Liu et al., 2024; Vara et al., 2019). As the accessibility of these approaches increases, so will our understanding of the role of chromatin architecture during meiosis and thus its role in the evolution of reproductive mechanisms (e.g., Álvarez-González & Ruiz-Herrera, 2025; Chen et al., 2020).



**Figure 1.** Summary of reproductive strategies literature and high-quality genome assemblies across eukaryotes. (A) Published primary literature by phylum and year. Literature libraries were searched using Dimensions (on 8th May 2024), with a search prompt that included 36 terms relating to topics within reproductive strategies research (search prompt: hermaphrodite OR monoecy OR dioecy OR gynodioecy OR androdioecy OR gynomonoecy OR andromonoecy OR polygamodioecy OR polygamomonoecy OR apomictic OR gonochorous OR parthenogenetic OR sex chromosome OR sex determination OR arrhenotoky OR haplodiploid OR genome elimination OR hybridogenesis OR dosage compensation OR self compatibility OR self incompatibility OR hermaphroditism OR monoecious OR dioecious OR gynodioecious OR androdioecious OR gynomonoecious OR andromonoecious

In summary, methodological advances like those listed above promise exciting opportunities for closing existing gaps in our knowledge of reproduction. The Tree of Sex consortium is thus committed to accommodating any such data type that could be pertinent to the interpretation of reproductive diversity to facilitate a holistic understanding of reproduction.

## Comparative analyses and their role in advancing the field of reproductive strategy evolution

Centralizing and integrating information across the field will enable large, phylogenetically deep comparative analyses to test for the evolutionary forces that shape diversity in reproductive strategies. With such analyses we can estimate the rate of reproductive trait evolution, the likelihood of transitions from one strategy to another, test whether trait evolution or transitions correlate with environmental, ecological, physiological, or genetic changes, and reconstruct the reproductive strategies in the ancestors of extant taxa. By compiling information as exhaustively as possible across eukaryotes, the Tree of Sex database will facilitate analyses at a previously unfeasible scale and resolution. Below we highlight several broad topics and ways in which we envisage such comparative analyses might advance our knowledge of reproduction.

### The evolution of sexual vs asexual reproduction, and everything in between

**Sexual reproduction**, broadly viewed as a process that brings together and mixes the genetic material from two parental gametes, is widespread in eukaryotes (Speijer et al. 2015). At its core, sexual reproduction involves the production of haploid (reduced) cells via meiosis. These cells can either be gametes (in most animals), or spores that will develop and produce gametes after additional mitotic cell divisions (in land plants). Haploid gametes then fuse (**syngamy**) during fertilization to create the genome of the offspring. However, within this broad definition of sexual reproduction lies a large diversity of strategies (Figure 2). For example, sexual reproductive strategies can differ in relative time spent within individual stages of their cycle. In some cases, the reduced stage is dominant; in others, the unreduced stage is dominant, while others spend large portions of their life cycle between these stages in the heterokaryotic phase. Strategies also differ in the types of gametes or spores that they produce. Many sexual eukaryotes are **anisogamous**, producing two gamete types that differ in size and form (e.g., eggs and sperm in many animals), while others are **isogamous**, wherein all gametes produced are of similar size and form (e.g., brewer's yeast and some protists, dinoflagellates, and algae (Lehtonen et al., 2016)). A similar distinction can be made between spore-producing organisms

which can either make spores of similar size (i.e., isospory, found in bryophytes; Mogensén 1985 and most ferns; Dyer 1979) or different sizes (anisospory, as found in seed plants; Jesson & Garnock-Jones 2011). Lastly, in some species, syngamy occurs between the gametes from two individuals (biparental sexual reproduction); in others, zygotes can be made from the fusion of gametes from only one individual (uniparental sexual reproduction), and some species can do both.

The evolution of anisogamy has received much attention in the context of species with distinct gamete types carried by different individuals (Parker, 1978). Anisogamy is thought to be a major cause of the evolution of different sex phenotypes (Schärer et al. 2012), and of sexual selection (Janicke et al., 2016). But why does anisogamy evolve despite its tendency to restrict mating opportunities (Otto, 2009)? And why is the predominant number of gamete classes two (Charlesworth, 1978)? Many models have been proposed to answer these questions (Billiard et al., 2011), with some receiving theoretical support (Constable & Kokko, 2021; Lehtonen & Kokko, 2011; Lehtonen & Parker, 2014; Lehtonen et al., 2021). However, conclusive empirical evidence for any of them remains rare. One type of empirical evidence that could prove useful in this case is the association of gamete types with other fundamental reproductive traits (e.g., self-compatibility, sex ratios, gamete competition/fertilization success) though such analyses would require multiple phylogenetically independent taxa to provide statistical power. Thus, this topic could benefit greatly from the integrated knowledge of such traits across eukaryotes, and especially clades such as algae (Coelho et al., 2018; Krueger-Hadfield, 2024) and fungi (Billiard et al., 2011).

The two distinct types of gametes of anisogamous species can be produced by different individuals or within a single individual. When produced by different individuals, we refer to the species as **gonochoristic** in animals, or **dioecious** in plants with a diploid-dominant life cycle. While in plants with a haploid-dominant life cycle, species in which different gamete types are produced by separate gametophytes are termed **dioicous** (Villarreal and Renner 2013) (Figure 2A). If both gamete types are produced by a single individual, we refer to them as **hermaphroditic/monoecious/monoicous**. Hermaphrodites can further be classified into those in which individuals can produce both types of gametes at the same time (**simultaneous/synchronous hermaphroditism**) or at different times (**sequential hermaphroditism**). In plants, **hermaphroditism** refers to systems where individuals carry bi-sexual flowers, while the system where individuals generate male and female flowers is called **monoecious**. Hermaphroditism is the most common sexual system in flowering plants (Käfer et al., 2017; Renner 2014) and has been described in a variety of invertebrates (see initial review by Ghiselin, 1969); while among vertebrates,

OR polygamodioecious OR polygamomonoecious OR apomixis OR gonochoristic OR hermaphroditic OR parthenogenesis OR self compatible OR self incompatible). The resulting 85,533 papers were searched for taxonomic names taken from the NCBI taxonomy (downloaded 8th May 2024). A total of 60,409 could be classified into phyla (though no manual curation was performed, so some errors may exist). We included all name categories (e.g., scientific, common, blast names) for all taxonomic ranks between phylum and genus. Common names and blast names were also included for species, but species scientific names were not used to reduce the total number of search terms, and because genera were already included. Inset pie chart shows the total proportion of all classified papers by phylum, with four model organisms also shown. For comparison, the black line shows a 4.1% annual increase publication rate estimated as the rate for all scientific publications (Bornmann et al., 2021) (B) All genomes archived by the The International Nucleotide Sequence Database Collaboration by phylum and year (Created in GoaT (Challis et al., 2023) on 06 June 2024). Only one assembly per species is counted. (C) Estimate of the number of named species for the 15 most speciose phyla taken from the Global Biodiversity Information Facility as of 16th May 2024. (D) Distribution of sex chromosome information (morphology and system of heterogamety) from ToS v1 database across eukaryotic phyla (Created in GoaT from data sources here).

# Exemplar paths of reproduction

The diagram illustrates the life cycles of various organisms, categorized into two main phases: **reduced phase (haploid)** and **unreduced phase**.

**Reduced Phase (Haploid):**

- monokaryotic mycelium gametophyte:** Can undergo *vegetative reproduction via mitosis* or *mitosis* to become an **adult**.
- adult:** Can undergo *mitosis* to become a **spore** or *abortive meiosis* to become a **gamete<sup>1</sup>**.
- spore:** Can undergo *mitosis* to become a **gamete<sup>1</sup>** or *meiosis* to become a **germ-cell**.
- gamete<sup>1</sup>:** Can undergo *meiosis* to become a **germ-cell** or *syngamy<sup>2</sup> (plasmogamy)* to become a **dikaryotic mycelium**.

**Unreduced Phase:**

- germ-cell:** Can undergo *mitosis* to become a **zygote** or *parthenogenesis* to become an **unreduced gamete**.
- zygote:** Can undergo *mitosis* to become a **adult** or *meiosis* to become a **gamete<sup>1</sup>**.
- adult:** Can undergo *mitosis* to become a **sporophyte** or *vegetative reproduction via mitosis (budding, fission, runners ...)* to become a **monokaryotic mycelium gametophyte**.
- sporophyte:** Can undergo *meiosis* to become a **gamete<sup>1</sup>** or *syngamy<sup>2</sup> (karyogamy)* to become a **dikaryotic mycelium**.
- dikaryotic mycelium:** Can undergo *mitosis* to become a **basidium** or *syngamy<sup>2</sup> (karyogamy)* to become a **zygote**.
- basidium:** Can undergo *meiosis* to become a **gamete<sup>1</sup>** or *syngamy<sup>2</sup> (karyogamy)* to become a **zygote**.

**Legend:**

- canonical animal reproduction
- common processes in nature
- example taxa in their prominent phase

## 2 types of reproduction by the origin of gamete(s)

**1** types of fusing gametes

**anisogamy (unequal sizes)**

macrogamete (female gamete) microgamete (male gamete)

**isogamy (same size)**

mating types

**2** types of reproduction by the origin of gamete(s)

**biparental sexual reproduction amphimixis**

**uniparental reproduction**

**autogamy (selfing; syngamy of two gametes from the same parent)**

**parthenogenesis (no syngamy; reproduction of unfertilised eggs)**

Downloaded from <https://academic.oup.com/jeb/article/38/7/861/8126803> by Indiana University - Bloomington user on 04 August 2025

it is thought to be specific to teleost fishes (Pla et al., 2022). Hermaphrodites can also coexist with males (androdioecy), females (gynodioecy) or both (trioecy/trioicy), in mixed sexual systems. In animals (Sasson & Ryan, 2017; Weeks et al., 2006), androdioecy is known in very few taxa, for example, in *Caenorhabditis* nematodes (Thomas et al., 2012), in branchiopod crustaceans (Benvenuto & Weeks, 2020); in barnacles (Ewers-Saucedo et al., 2016) and among vertebrates, it has only been observed in *Kryptolebias* killifishes (Costa et al., 2010). Gynodioecy is even more rare in animals (Weeks, 2012). In flowering plants, gynodioecy is more common than androdioecy (McCauley & Bailey, 2009), while trioecy is very rare in both animals (Oyarzún et al., 2020) and flowering plants (Godin, 2022). Recent studies have investigated evolutionary transitions among all sexual systems (Goldberg et al., 2017; Leonard, 2019; Pannell & Jordan, 2022; Weeks, 2012). By centralizing all known examples of sexual systems and self-compatibility in eukaryotes, the Tree of Sex database will allow us to better test whether the contexts of these strategies are general across Eukaryota.

While sexual reproduction is often thought of as involving two individuals (biparental), in **self-compatible simultaneous hermaphrodites** the fusing gametes might be from the same parent in a process known as self-fertilization or selfing. Selfing is found in plants (Baker, 1955; Barrett, 1998; Hedrick, 1987) and animals (Goodwillie et al., 2005a; Jarne & Auld, 2006; Jarne & Charlesworth, 1993), in **automictic** fungi (Hood & Antonovics, 2004) and algae (Chepurnov et al., 2004; Heesch et al. 2021; Krueger-Hadfield et al., 2024). Uniparental sexual reproduction has been most extensively studied in the form of self-fertilization in plants (Cheptou, 2018; Hartfield et al., 2017; Wright et al., 2013), which is often thought to evolve in response to selection for reproductive assurance when mates (or pollinators) are scarce (Baker, 1955; Pannell et al., 2015). Selfing in animals has been reported in a variety of invertebrates (Annelida, Arthropoda, Cnidaria, Echinodermata, Ectoprocta, Mollusca, Nematoda, Platyhelminthes, and Urochordata; Jarne & Auld, 2006). Isogamous species might undergo self-fertilization via fusion of gamete of the same meiotic tetrad (Billiard et al., 2011). Unfortunately, many selfing taxa, e.g., algae, still lack population genetic data with which to test theories for the predictors and consequences of selfing (Krueger-Hadfield, 2024). However, by centralizing known examples of gonochorism, hermaphroditism, and self-fertilization across diverse lineages, the Tree of Sex database will allow us to search for explanations for the distribution of selfing across eukaryotes.

Uniparental reproduction might also happen without fusion of gametes (syngamy), which is known as **parthenogenesis**. These species usually have a modified meiosis or pre- or post-meiotic genome duplication, resulting in unreduced gametes and thus to stable ploidy levels between generations (Stöck et al., 2021b). The offspring might be a genetic clone of the parent (e.g., in parthenogenetic animals using premeiotic endoreplication; Dedukh et al., 2024) or a mixture of parental haplotypes known in plant and animal literature as **automixis**. Finally, reproduction can occur via separation of somatic tissue such as vegetative propagation in plants (Schwaegerle, 2005), budding or fragmentation in cnidarians (Berzins et al., 2021), fissiparity in some flatworms (Zaccanti & Farne, 1986) and echinoderms (Dolmatov et al., 2018; Thuy et al., 2024).

Though we have presented sexual and asexual reproduction above as separate modes of reproduction, in fact a spectrum of strategies exists, which do not neatly fall into these two discrete categories. For example, in sperm-dependent parthenogenesis, also known as gynogenesis, sperm is required but only to initiate embryo development, i.e., its DNA is not incorporated into the zygote (Beukeboom & Vrijenhoek, 1998). This system is known, for example, in the teleost fish genera *Poecilia* and *Poeciliopsis* (Cerepaka & Schlupp, 2023; Lampert & Scharl, 2008; Schlupp, 2005; Schultz, 1969), *Carassius* (Choleva et al., 2008; Komen & Thorgaard, 2007; Liet al., 2018) and *Cobitis* (Choleva & Janko, 2013). Similarly, androgenesis requires two parents for fertilization, but a zygote is formed solely with paternal nuclear genes; this is found in plants and animals (Schwander & Oldroyd, 2016). Some amphibians, teleosts and insects perform hybridogenesis, wherein DNA from both sperm and egg are incorporated into the zygote; however, the genome of one of the parental species is consistently eliminated during meiosis in each generation (Neaves & Baumann, 2011).

The categorization of organisms as exclusively sexual or asexual also does not work for organisms using more than one reproductive strategy, as in cyclical parthenogenesis (found in aphids and water fleas) or **arrhenotoky** (a form of haplodiploidy) where males develop parthenogenetically from unfertilized eggs and females develop sexually from fertilized egg (for example in hymenopterans, thrips, and several other arthropod lineages; reviewed in de la Filia et al., 2015). Finally, many species switch between sexual and asexual forms of reproduction. Such strategies are used by most studied unicellular eukaryotes which generally reproduce asexually, with occasional sexual cycles (Green & Noakes, 1995; Hofstatter & Lahr, 2019).

Obligately asexual (or uniparental) reproduction is relatively rare, which has led to the famous question “why sex?” (Bell, 1981; Hartfield & Keightley, 2012; Otto, 2009, 2021). Indeed, where, how, and why sexual reproduction and its constituent processes (meiotic recombination, outcrossing, anisogamy, and sex) have evolved or been lost has received much attention. It is thought that recombination dates back to the ancestor of all eukaryotes (Bernstein & Bernstein, 2010; Mirzaghaderi & Hörandl, 2016; Ramesh et al., 2005). Yet, recombination has been lost or much reduced in the meiosis of the heterogametic sex in many species (Morgan, 1914; Sardell & Kirkpatrick, 2020; Satomura et al., 2019), or lost entirely in some asexuals employing asynaptic meiosis, like the Amazon molly (Dedukh et al., 2022) or achiasmatic meiosis, as in *Bithynia* snails (Debus, 1978) and scorpions (Shanahan & Hayman, 1990). It has been proposed that recombination evolved because it disrupts detrimental combinations of alleles and brings together beneficial alleles that arise in different individuals (Felsenstein, 1974; Hartfield & Keightley, 2012; Otto, 2009). Furthermore, recent theoretical work points to selective interference as the main driver of the evolution of sexual reproduction with recombination (Otto, 2021). Without genetic mixing and recombination, selection cannot act upon individual loci, as genetic variance is restricted to the level of the whole genome. There is substantial empirical support for selective interference (Neiman et al., 2018); however, so far it is restricted to individual genes and limited taxonomically. Importantly, many of these hypotheses have remained largely untested, primarily due to technical challenges in accurately estimating key parameters, such

recombination rates, effective population sizes, and selection coefficients (Dapper & Payseur, 2017), as well as the fact that sex and meiotic recombination evolved only a small number of times. Thus, there is yet to be a fully satisfactory explanation for the vast prevalence of sex, which necessarily needs to be based on a study of a taxonomically diverse set of asexual species.

The complexities of categorizing reproductive strategies make comparative analyses difficult. By structuring the database to include nuanced characteristics of meiosis and inheritance for these species, the Tree of Sex will allow us to study the evolution of various components of these reproductive strategies in their most relevant context without forcing them into categories that could mask important patterns.

### Explaining the many ways of sex determination

Anisogamy is more common than isogamy among multicellular eukaryotes and is often associated with dioecy/dioicy/gonochorism. In such species, mechanisms must exist to determine which type(s) of gamete an individual will produce, and individuals are classified as female (producing macrogametes) or male (producing microgametes). We refer to such mechanisms as sex determination (Beukeboom & Perrin, 2014; Bull, 1983). This term is also sometimes used in isogamous clades such as algae and fungi, though, to avoid confusion, we refer to individuals producing different types of similarly sized gametes as **mating types** (Aanen et al., 2016).

One of the main goals of the first Tree of Sex project was to synthesize the many ways in which sex is determined across eukaryotes and the non-random distribution of these mechanisms among eukaryotic clades (Bachtrog et al., 2014). For instance, to our knowledge every mammal or bird has **genotypic sex determination** (GSD), wherein the sex of an individual is determined by their genotype at one or more locus, or in their karyotype. In contrast, some reptiles and teleost fishes use **environmental sex determination** (ESD), wherein male or female development depends on environmental cues (e.g., temperature or social environment).

Similarly to reproductive strategies, however, sex determination modes resist binary classification into GSD or ESD with sex in some species being determined by the interaction between genotype and the environment (Baroiller et al., 1995; Collin, 2006; Holleley et al., 2015; Mork et al., 2014; Piferrer et al., 2005; Sarre et al., 2004; Whiteley et al., 2021). In addition, there are increasing examples of putative GSD systems in which the genotype at the presumed sex determination locus is less predictive of the sex phenotype than originally thought (Cossard & Pannell, 2019; Ehlers & Bataillon, 2007; Holleley et al., 2016; Nemesházi & Bókony, 2022; Nemesházi et al., 2020; Phillips et al., 2020; Rodrigues et al., 2018; Wiggins et al., 2020). It has been proposed (but not yet empirically tested) that such cases could arise from interactions between the random noise inherent in gene expression and the required expression thresholds for sex development pathways (Perrin, 2016). Though, it is also possible that the occurrence of mismatches between genotypic and phenotypic sex could be favoured by natural selection as it can allow for occasional self-fertilization, which could be beneficial in some situations (e.g., low mate abundance, Crossman and Charlesworth 2014; Pannell 2015).

There is also ample variation in sex determination mechanisms within GSD systems. In many cases a single locus acts, in a switch-like fashion, to divert development to the

male or female pathway. This gene is widely referred to as the Master Sex Determination gene, though some suggest that it should be referred to as the Primary Signal (PS) gene reflecting a more polygenic view of developmental regulation (Kocher et al., 2024). We use the latter from hereon. Polygenic sex determination has indeed been shown in some species, such that multiple genes in an individual's genome act together to determine sex (reviewed in Kocher et al., 2024; Schartl et al., 2023). Another form of a GSD present in several diverse taxa (15% of known arthropod species; de la Filia et al., 2015) is arrhenotoky, which has independently evolved multiple times (Normark, 2003; Ross et al., 2015) and is often associated with eusociality (da Silva, 2022; Joshi & Wiens, 2023), inbreeding (Hamilton, 1967; Ross et al., 2015), flexible sex allocation (Hamilton, 1967; Normark, 2004), reproductive manipulation by infectious endosymbionts (Werren et al., 2008) and often uses the ecologically vulnerable complementary sex determination system (Beye et al., 2003; Leung & van der Meulen, 2022; Zayed & Packer, 2005).

Despite great progress in the discovery of sex determination mechanisms across eukaryotes (Bachtrog et al., 2014; Beukeboom & Perrin, 2014; Blackmon et al., 2015, 2017; Goldberg et al., 2017; Iwasaki et al., 2021; Sabath et al., 2016), this information is distributed in a very biased way across eukaryotic clades. For example, we currently know the PS gene of approximately 140 teleost species (Kitano et al., 2024; Wang et al., 2024), which has allowed us to recognize not only the high amounts of polymorphism in PS genes in this clade, but also the strong convergence on a few genes, implying strong evolutionary constraints on which genes can become PS genes in some lineages. These insights highlight how important and powerful it is to have good phylogenetic coverage and detailed information on reproductive strategies. In contrast, we currently have no knowledge of the sex determination genes in most organisms, including even well-studied ones like guppies (Charlesworth et al., 2020) or *Rumex* plants (Sacchi et al., 2024) to name but two cases. This not only precludes similar inferences in these clades but also prevents our understanding of whether the evolutionary patterns seen within teleosts are an exception, or whether they are indicative of a general pattern across eukaryotes. This issue applies not only to the study of PS genes, but also to mechanisms of ESD, which, to our knowledge have only been described to date in turtles (Ge et al., 2018; Schroeder et al., 2016; Weber et al., 2020) and *Daphnia* (Kato et al., 2024), though similar ESD mechanisms involving thermally sensitive isoforms have been implicated across distantly related reptiles (Deveson et al., 2017; Whiteley et al., 2022).

By integrating available data on sex determination systems across eukaryotes, the Tree of Sex database will facilitate comparative approaches to describe, with higher resolution than before, the distribution of sex determination mechanisms across eukaryotes, their association with relevant traits (e.g., life cycle, environmental stability), and patterns of gene use for sex determination.

### Sex chromosome evolution

In organisms with GSD, the term sex chromosome refers to a linkage group harbouring a locus whose inheritance correlates with an organism's sex. They are common among eukaryotes with separate sexes and have evolved many times independently across many clades (Bachtrog et al., 2011; reviewed in Bachtrog et al., 2014). The term "sex chromosome" refers

to all ranges of sex chromosome differentiation (see below), including systems in which one homolog is almost completely sex linked, completely absent, or in which the sex-linked region containing the sex determination gene represents only a small portion of the chromosome, with the rest segregating autosomally (**pseudoautosomal region [PAR]**). In organisms with fixed separate sexes, in which sex is determined in the diploid phase of the life cycle, sex chromosome systems can be split into two major categories: (1) male **heterogamety** (e.g., XX/XY), where males (XY) can produce two different (hence “hetero”) types of haploid gametes, containing either an X or a Y chromosome; and (2) female **heterogamety** (e.g., ZZ/ZW), where it is the females that produce two different gamete types (e.g., with either a Z or a W chromosome). In such systems, the sex-limited chromosome may carry an allele that initiates development of the heterogametic sex, or sex might be determined by the number of X (Bridges, 1916; Meyer, 2022) or Z (Ioannidis et al., 2021) chromosomes. In some organisms, no sex-limited chromosome exists (e.g., XX/X0 in nematodes (Nigon, 1949), springtails (Núñez, 1962), and many insects (reviewed in Blackmon et al., 2017; White, 1945), or ZZ/Z0 in many moths (Sahara et al., 2012)). Again, in such cases, it is the number of X or Z chromosomes that determines sex. And finally, taxa in which *only* the sex-limited chromosome exists have also been found (00/Y0, or 00/W0) (Jonika et al., 2022), though the latter is seemingly rare. In anisogamous species in which sexes are determined in the haploid life phase, sex chromosomes are denoted as U/V, with only one of the two being present in a given haploid cell (Ahmed et al., 2014; Lipinska et al., 2024; Nieuwenhuis & James, 2016). Finally, while isogamous species do not have sex chromosomes, their “mating-type chromosomes” often behave similarly to U/V chromosomes (Coelho et al., 2018).

One of the most notable and consistent properties of sex chromosomes is the tendency for recombination to be reduced between the sex-limited chromosome (e.g., Y or W) and its **gametolog** (e.g., X or Z) (Charlesworth, 2023; reviewed in Charlesworth et al., 2005). This can either be a progressive reduction (as reviewed in Jay et al., 2024), or perhaps due to a sex determination gene arising in a region that already lacked recombination (Bergero et al., 2019; Rifkin et al., 2020, 2022). Due to the lack of genetic exchange between gametologs, the sex-linked region often degenerates over time, losing much of its gene function and content (Muller, 1964) and accumulating repetitive elements (Charlesworth et al., 1994; Nguyen & Bachtrog, 2021; Peona et al., 2021). However, due to differences in selection in the two sexes, sex chromosomes can also become specialized, e.g., Y chromosomes may become enriched in genes important for male reproduction (Bachtrog et al., 2019; Carvalho et al., 2001; Colaco & Modi, 2018; Okada et al., 2001). In contrast to XY/ZW systems, the degeneration and/or specialization of U and V chromosomes is expected to be symmetrical as they are each inherited without recombination by gametophytes of just one sex (Bull, 1978; Carey et al., 2021; Immler & Otto, 2015).

The evolutionary forces driving recombination reduction on sex chromosomes have received considerable debate. Several theoretical models have been proposed to explain this process (reviewed in Charlesworth, 2023; Jay et al., 2024; Käfer, 2022; Ponnikas et al., 2018; Veltsos et al., 2024). Most of these theories are adaptive (but see Jeffries et al., 2021; Kent et al., 2017; Rifkin et al., 2020) and invoke selection either due to the presence of sexually antagonistic

polymorphisms, deleterious mutations, regulatory evolution, or meiotic drive. While debate is ongoing as to which of these models is most promising (Bergero & Charlesworth, 2019; Charlesworth & Olito, 2024; Jay et al., 2024; Lenormand & Roze, 2024), this discussion is almost entirely theoretical and rests on many untested assumptions (summarized in Jay et al., 2024). However, some of these assumptions can potentially be tested (Charlesworth, 2023) and the Tree of Sex database will facilitate this, for example, by relating the rate and extent of recombination loss on sex chromosomes to its proposed drivers across phylogenetically distant species.

Another interesting characteristic of sex chromosomes is the rate at which fusions between sex chromosomes and autosomes fix within some lineages (Anderson et al., 2020; Lisachov et al., 2021; Pennell et al., 2015; Pokorná et al., 2014; Wright et al., 2024). This is often followed again by the spread of recombination reduction to parts of the fused chromosome (**neo-sex chromosome**), further increasing the proportion of the genome which is sex linked. In extreme cases, this can happen multiple times, as in the sex chromosome chains of monotremes (Rens et al., 2007). Proposed evolutionary drivers of such fusions overlap with those proposed above for recombination suppression. For example, a sex chromosome-autosome fusion may be favoured if the autosome harbours a gene under sexually antagonistic selection (Charlesworth & Charlesworth, 1980; Matsumoto & Kitano, 2016). However, additional hypotheses specific to the fixation of fusions also exist. The fragile Y hypothesis (Blackmon & Demuth, 2015b), for example, proposes that, as recombination is lost on a sex chromosome, opportunity for chiasmata is reduced along much of its length. As chiasmata are important for correct chromosomal segregation during meiosis, their absence can result in maladaptive or lethal aneuploidies. This could provide an advantage to sex chromosomes fused to an autosome, which could restore pairing. Here, the Tree of Sex database will facilitate a more complete characterization of the rates of various types of fusions across the eukaryotic phylogeny. Further, it will allow for tests of association between the rate of fusions and the extent of sex chromosome degeneration, size of the pseudoautosomal region, the integrity of centromeres, holocentricity, and many more potential predictors across a much more diverse set of lineages than is currently possible.

Extensive degeneration via deletions, massive accumulation of repeat elements (Chalopin et al., 2015), or fusions can all result in **heteromorphic** sex chromosomes, i.e., those that show morphological differences that can be seen via microscopy. As such techniques were the only ones available for decades after sex chromosomes were first discovered in 1905 (Brush, 1978; Stevens, 1905, 1906), heteromorphic sex chromosomes were generally the only ones that could be detected and studied. However, the increasing availability of genetic and genomic techniques has enabled the discovery of many new sex chromosome systems that do not show morphological differences via microscopy (**homomorphic**). While this newly available data is still fragmented, it has revealed that homomorphic sex chromosomes are common among eukaryotes (Bachtrog et al., 2014). Further, the distribution of homomorphic vs heteromorphic sex chromosomes across eukaryotic lineages is highly heterogeneous. For example, in mammals almost every species has heteromorphic sex chromosomes (Graves, 1995), while in amphibians, between 75% and 96% of species are thought to have homomorphic sex

chromosomes (Ma & Veltsos, 2021; Schmid et al., 1991). Assembling detailed information for sex chromosome differentiation from all data types, including microscopy, genetic maps and genomic data within the Tree of Sex database will help crystallize this picture, and give a much more accurate view of the extent and distribution of sex chromosome degeneration across the eukaryote phylogeny.

The extent of sex chromosome degeneration observed is related to the frequency at which new sex chromosomes arise in each clade. Transitions from hermaphroditism to gonochorism (with GSD; Ming et al., 2011), ESD to GSD (Johnson Pokorná & Kratochvíl, 2016; Sabath et al., 2016), or from one GSD system to another (i.e., sex chromosome turnovers; El Taher et al., 2021; Gamble et al., 2015; Jeffries et al., 2018; Kratochvíl et al., 2021b; Vicoso, 2019) will all create new sex chromosomes. Because, in most cases, new sex chromosomes arise from existing autosomes (but see Fraïsse et al., 2017; Imarazene et al., 2021), and because sex chromosomes degenerate progressively through time, the gametologs of very young sex chromosome systems are generally undifferentiated. Lineages with frequent transitions in sex determination system thus tend to have undifferentiated sex chromosomes (Blaser et al., 2014; El Taher et al., 2021; Jeffries et al., 2018). However, correlations between the age of a sex chromosome system and levels of degeneration can be weak across divergent taxa (Renner & Müller, 2021) and there are several known cases of old yet homomorphic sex chromosome systems (Kuhl et al., 2021; Pan et al., 2019). Given the ratchet-like nature of sex chromosome degeneration, this may reflect differences in the rates of sex chromosome degeneration, but we currently lack the dataset to assess this. To solve this question, we need accurate estimates of the age of sex chromosomes (e.g., from phylogenies and neutral sequence divergence (dS)) along with accurate measures of their degeneration (e.g., gene loss, dNdS, repeat accumulation) from multiple species within multiple distant clades. Indeed, such a task is a prime candidate for the new Tree of Sex database and consortium.

Another mystery relating to sex determination system transitions is their highly heterogeneous rates among taxa. For example, they are very rare in some lineages within insects (Toups & Vicoso, 2023), bryophytes (Carey et al., 2021), cephalopods (Coffing et al., 2024), sturgeons (Kuhl et al., 2021), birds (Fridolfsson et al., 1998) and many other amniote lineages (Kratochvíl et al., 2021a), but very frequent in geckos (Gamble et al., 2015), true frogs (Jeffries et al., 2018), medaka fishes (Ansai et al., 2022; Myosho et al., 2015), and cichlids (El Taher et al., 2021; Feller et al., 2021). Why some sex chromosome systems are so stable, while others transition often, remains an important but unresolved question. Comparative analyses are essential for identifying, dating, and placing transitions on a phylogeny. The Tree of Sex will allow us to do this on an unprecedented scale, and by relating transitions to other life-history traits, we hope to better understand their evolutionary drivers.

### Evolutionary and ecological consequences of reproductive strategies

The reproductive strategy of a given taxon affects its ecology and evolution. For instance, dispersal to, and establishment in a new geographic region can be easier for self-compatible hermaphroditic or asexual species, especially in geographic regions where pollinators or mates are rare (Baker, 1955). This theory has received support in many hermaphroditic

plant systems (Grossenbacher et al., 2015; Hargreaves & Eckert, 2014; Pannell et al., 2015; Razanajatovo et al., 2016). Similarly, in animals, simultaneous hermaphroditism can increase fertilization success in unpredictable, unstable, or extreme environments where mates are at low density (Benvenuto & Weeks, 2020), as for parasitic species, and gynogenesis using sperm from other species can result in increased invasion success where conspecific mates are not present (Fuad et al., 2021). For example, parthenogenesis has been linked to range expansion in stick insects (Morgan-Richards et al., 2010). Extending this test of range size to other modes of reproduction would give a more holistic picture of how colonization and establishment can be impacted by reproductive strategy (Tilquin & Kokko, 2016). This could further be used to help predict the potential invasiveness or pathogenicity of species, or their potential for range shifts in response to climate change or other anthropogenic impacts.

Reproductive strategies also influence the ability of taxa to adapt and thus their resilience to biotic or abiotic environmental change. The efficacy of selection is impacted by random associations that occur between alleles that increase fitness and those that decrease fitness limiting both adaptation and the elimination of deleterious mutations (reviewed in Otto, 2021). This process is known as Hill–Robertson effect (Hill & Robertson, 1966) or “selective interference.” Sexual reproduction with recombination can break down these genetic associations and generate variation upon which selection can act. However, some reproductive strategies, such as self-fertilization, limit the effective rate of recombination and therefore exacerbate selective interference (Hartfield, 2016; Hartfield et al., 2017). Similar consequences are predicted in parthenogenetic species, though the exact dynamics depend on the cellular mechanism of parthenogenesis (Engelstädter, 2017). Furthermore, the origin of parthenogenetic species is another factor affecting the levels of selective interference. In addition, the efficacy of selection can be reduced when new parthenogenetic or selfing species form, if they arise spontaneously with little genetic diversity (Jaron et al., 2022).

In the absence of sexual reproduction and effective recombination in a diploid species, homologous haplotypes will independently accumulate mutations resulting in extreme Allelic Sequence Divergence (White, 1945), often also referred to as the “Meselson effect” (Birky, 1996). Consequently, corresponding haplotypes in different individuals and/or populations will be more closely related than two homologous haplotypes within an individual. Despite the elegance of this predicted effect and a large number of parthenogenetic species tested, support is limited (Brandt et al., 2021; Öztoprak et al., 2023; Weir et al., 2016) and evidence in several systems has later been reinterpreted (Freitas et al., 2023; Jaron et al., 2022; Mark Welch et al., 2008; Schwander et al., 2011). Hybridization of distantly related species can generate new parthenogenetic lineages that also show a similar genomic pattern to the Meselson effect (Jaron et al., 2021). Some parthenogenetic species occasionally lose heterozygosity by mitotic recombination between homologous parental chromosomes (Janko et al., 2021), which potentially contributes to mitigation of the accumulation of deleterious mutations due to selective interference (Kočí et al., 2020). In general, the consequences of parthenogenesis have proven less apparent compared to early predictions (Normark, 2003) and are often masked by lineage-specific effects (Jaron et al., 2021). It is of great interest to separate lineage-specific effects and the

consequences of the individual reproductive strategies. Only then can we identify the underlying causes of the observed phenomena and finally get closer to answering the question “why sex?”

In the face of escalating environmental change, determining how different reproductive strategies interact with the environment to influence population resilience is also key for conservation biology. Of the environmental stimuli that can influence organisms, temperature can affect sex determination in many vertebrates (Kitano et al., 2024; Nemesházi & Bókonyi, 2022; Valenzuela & Lance, 2004). The fast pace of global warming threatens temperature-sex determined vertebrates by disrupting sex ratios and other traits (Honeycutt et al., 2019; Jensen et al., 2018; Lockley & Eizaguirre, 2021; Valenzuela et al., 2019). Similar climate-driven sex-ratio distortions are expected in GSD species that are prone to thermal sex reversal, with complex consequences for population dynamics and microevolution (Bókonyi et al., 2017; Nemesházi et al., 2021; Schwanz et al., 2020; Whiteley et al., 2018). Further, population dynamics of species with multifactorial sex determination such as GSD houseflies or TSD/GSD silverside fish, where the distribution of sex determination variants is temperature-dependent and geographically clinal (Duffy et al., 2015; Feldmeyer et al., 2008; Foy et al., 2024), will also be affected by climate change. Aside from temperature, sex determination mechanisms may also be disrupted by pollutants that act as endocrine disruptors that directly affect sexual development or induce stress that renders individuals susceptible to thermal insults (Mizoguchi & Valenzuela, 2016). Thus, anthropogenic disruption of the chemical (pollution) and thermal environment (globally or locally, e.g., the urban heat island effect) may have a combined influence on sex determination. For instance, elevated sex-reversal frequency has been observed in agile frogs (*Rana dalmatina*) in areas where anthropogenic land use (i.e., agriculture and urbanization) is high (Nemesházi et al., 2020). By centralizing data for environmental effects on sex determination mechanisms, the Tree of Sex database will allow a comprehensive overview of species that are more vulnerable to unavoidable environmental changes, which could, in turn, support conservation actions.

At a macroevolutionary scale, reproductive strategies play an important role in shaping species diversity. For instance, in the plant family Solanaceae, obligate outcrossing (self-incompatibility) is associated with higher net diversification rates than in species capable of self-fertilization (Goldberg et al., 2010). Even though self-compatible Solanaceae species exhibit a higher speciation rate than obligate outcrossing species, this rate is exceeded by the much higher extinction rate of self-compatible plants. This pattern agrees with theoretical expectations, showing that species barriers accumulate faster in selfing than outcrossing species under most scenarios of speciation (Marie-Orleach et al., 2022). However, the relative importance of genetic (selective interference) and demographic factors in driving higher rates of extinction in self-compatible species remains unclear. Species with separate sexes can exhibit conflicts for traits that have different optima between sexes, leading to an evolutionary arms race that can ultimately contribute to reproductive isolation. A comparative analysis of hybridizing plants supports this idea, with strictly dioecious species pairs exhibiting an excess of species barriers compared to species with other reproductive strategies (Pickup et al., 2019). Sex chromosomes are also expected to

facilitate reproductive isolation, as evidenced by the two rules of speciation, namely Haldane’s rule (Haldane, 1922) and the large-X effect (Charlesworth et al., 1987; Coyne, 1985, 2018; Masly & Presgraves, 2007; Turelli & Orr, 1995; Turner & Harr, 2014). These rules have been widely validated using empirical data (reviewed in Beukeboom & Perrin, 2014). Furthermore, hemizyosity of sex chromosomes appears to be a key feature in explaining the disproportionate role of sex chromosomes in speciation (Fraïsse & Sachdeva, 2021; Lima, 2014); however, more work is required to understand the underlying mechanisms. It is also not known what impact sex chromosome turnovers have on reproductive barriers, though many potential mechanisms exist for them to do so. Using the Tree of Sex to relate these factors to rates of branch splitting on a robust phylogeny could lend valuable insights into the role of reproductive strategies in diversification.

Finally, the fundamental transition between isogamy and anisogamy is likely to have had dramatic consequences for sexual trait evolution. It is proposed to have caused stronger sexual selection in males leading to female-biased parental care and male-biased sexual dimorphism (Janicke et al., 2016). Mating systems also influence the costs and benefits of parental care for both sexes. For example, theoretical models predict that males increase the level of care they provide to their offspring when the certainty of their paternity increases (Westneat & Craig Sargent, 1996). However, the empirical evidence for a positive relationship between certainty of paternity and paternal care is mixed (Alonzo, 2010), which calls for more theoretical work and standardized empirical approaches.

In summary, the centralization and integration of reproductive knowledge in the Tree of Sex will allow researchers to address many fundamental questions related to the evolution of reproductive strategies. However, there are many more questions that could be addressed than those listed above. As the database will be a long-standing public resource, our hope is that the wider community will eventually use it to address questions that we have not yet conceived.

## The Tree of Sex rebooted

This new iteration of the Tree of Sex was established in June 2023, almost a decade after the first Tree of Sex project ended. Motivated by the questions and challenges above, we have two major goals: (1) to *centralize and integrate* published knowledge related to reproductive biology in a single public database and (2) to *provide a framework for ethical research and collaboration* in which to use the database to address scientific questions. Below we outline our plans, current progress and the challenges we foresee for the Tree of Sex consortium and for the database implementation.

## The consortium

At the time of writing (April 2025), the consortium has approx. 200 members, from 130 institutions across 35 countries. The Tree of Sex consortium is a global community of scientists with the aim of building a collective knowledge base in the field of reproductive biology of eukaryotes. We are currently in the process of establishing the foundations for a long-term, self-sustaining initiative (Figure 3). To accomplish this task, the consortium is managed by a general committee and six “operations teams,” each responsible for specific demands of the consortium (Table 2). These demands are

currently focused on setting up the infrastructure necessary for the development of the database in a sustainable and equitable manner.


Given the volume of literature, numerous subdisciplines and data types, and the great taxonomic breadth we aim to capture, we necessarily depend on the engagement of many experts. The Tree of Sex project is thus, by necessity, a community effort. The consortium is open, and we invite anyone and everyone interested in reproductive biology to join us via the registration form at the Tree of Sex webpage (treeofsex.org). We welcome and encourage researchers from all career levels, nationalities, cultures, genders, and sexual orientations, especially those which have traditionally been minoritized in science.

Consortium challenges

Establishing a new consortium comes with several challenges, many of which are shared across projects. Here we will discuss two challenges that specifically concern the Tree of Sex. Firstly, reproduction is an extremely complex biological process but one that is always viewed through a lens of human bias; see [de Vries and Lehtonen \(2023\)](#) and references therein. Our aim within the Tree of Sex is to capture this complexity in a way that is as objective as possible. But how do we accomplish this when recognizing one’s own biases, or those of one’s group, is extremely difficult? We believe that the Tree of Sex consortium *must* equitably include a diversity of people from the start, so that they can co-determine

its scope, and recognize each other’s biases. To help accomplish this, the consortium includes a team dedicated to Ethics and Inclusivity ([Table 2](#)) whose specific goals centre around building a diverse and inclusive community of scholars. So far, the Ethics and Inclusivity team have laid the foundation for an inclusive and supportive environment for its members through the creation of a code of conduct for members of the consortium and attendees of networking events as well as an ethical vision to be used as a guide for ethical behaviour across all consortium activities. The team is currently working on establishing standard protocols for accreditation of work and systems to initiate discrete projects and allocate leadership and authorship opportunities in an equitable way. The future plans for this team include the tracking of recruitment, retainment, and advancement of consortium members from different backgrounds in order to monitor equitable inclusion and barriers to participation. We also envisage this team collaborating with others in the consortium to organize activities that mitigate inequity among members. Such activities may include, for example, bursaries for travel and childcare, mentorship programmes for students and members of minority groups within the consortium, or facilitating remote access to computing resources required to work with the ever-increasing amounts of genomic data that is relevant to much of the research on reproduction. Global inclusivity is a major outstanding issue in science, and we are committed to working towards an equitable consortium, yet we recognize the challenge in doing this effectively and we will continuously update our activities to do so.

The second challenge is concerning the socially charged nature of the topic of sex. Indeed, terms like sex, sexuality, and gender have a long history of critique from the feminist and LGBTQIA+ groups for their use to marginalize members of these communities ([Aghi et al., 2024](#); [Fausto-Sterling, 2008](#); [Richardson, 2013](#)). While the scientific work of the consortium will primarily focus on biological systems that have very little to do with the human-centric sociocultural context of sex, arguments using natural systems are frequently featured in public discussions. Therefore, we must put measures in place to prevent misuse of the collected data in pseudoscience and bigoted narratives. The exact steps we take to do this are a matter of ongoing debate within the consortium, and we welcome input from all members of the scientific community. Interpretations of sex or sexuality can vary widely even among biologists, and we are thus working hard to



**Figure 3.** Time plan for the three phases of the Tree of Sex consortium. We are currently advancing in setting up the infrastructure to manage the consortium and create the database. The next phase, starting this year, will be to fill the database with records and continue to keep the database updated. Once the database contains an adequate number of records, we will start analyzing the data by organized topic research teams.

**Table 2.** Operations teams currently active in the ToS Consortium.

Operations team	Deliverables
Database integration	- listing and describing reproductively relevant databases - facilitating their integration within Tree the Sex Database
Ethics and inclusivity	- guiding and formulating shared values of the consortium towards building a diverse community of scholars and approaching the topics of sex with awareness and sensitivity to its potential societal impact - supporting teams with the work of building the database and challenging decisions that may arise with our values of justice, equity, and inclusion
Funding	- monitoring research funding landscape for opportunities for the consortium and consortium members - supporting individual grant applications of consortium members
Literature search	- generating a list of relevant literature to be processed for the database - exploring opportunities of automated literature processing
Ontology	- create and maintain the Tree of Sex Ontology
Outreach and science communication	- internal communications of the consortium - managing social media and outreach events

construct a consortium and an environment where multiple interpretations of sex and related topics can be integrated. We also hope to collaborate with social scientists to discuss how our understanding of reproduction from nature feeds into human societies. And lastly, we are committed to tracking the press arising from the outputs of the consortium, and, where needed, writing public press pieces to counter any erroneous arguments directly based on our work.

### The database

The new Tree of Sex database will store the data at the level of observation (rather than species). This will allow for multiple observations of the same feature (e.g., chromosome numbers) for each species and will enable us to capture conflicting data within the literature regardless of whether they reflect erroneous results or biological polymorphism (e.g., in cases where individuals or populations differ in a reproduction-related trait). The database interface will allow users to aggregate data in multiple ways, including by taxon. For each piece of information stored in the database, the taxon will be recorded in the form of a taxon ID (TaxID) maintained within the International Nucleotide Sequence Database Collaboration. TaxIDs are unique and permanent identifiers for species that aggregate synonyms as well as accommodate species name changes. New identifiers will be requested following the guidelines of NCBI together with the Biodiversity Genomics community (Blaxter et al., 2024b). Each submitted observation will require a reference to its primary source (e.g., scientific article), with multiple submissions from the same reference possible. Finally, each record will be associated with the recorder via their ORCID (required for data submission), and the date of the data entry. These metadata ensure traceability of information and enable crediting active members of the community.

### Database challenges

The scope and scale of this new version of the Tree of Sex database present several challenges. First, the volume of research on reproduction is vast. The literature search performed to create Figure 1, though based on a limited search prompt, resulted in over 85,000 publications, which is likely the tip of the iceberg for information relevant to the database. Thus, while many papers may be redundant (e.g., thousands of papers on humans and mice), the relevant literature is enormous and growing at a cumulative annual rate of over 4% (Figure 1). Synthesizing this vast and growing literature is a daunting task, even with a large group of volunteer researchers. However, preliminary tests show that use of Large Language Models (LLMs) could greatly speed up this process. LLM tools can be used to interrogate publication PDFs and, with well-constructed prompts, extract specific pieces of information. Our current plans involve integrating these tools in a data extraction pipeline, whereby LLMs produce targeted information summaries of each paper, which are formatted to allow for automatic import into the database if deemed accurate. Experts will be tasked with developing these pipelines and checking summaries against the original paper. The final goal is to construct a hybrid AI-human pipeline in which (ideally) most papers can be automatically imported via the LLM-produced summary, with a proportion (e.g., 10%) being checked by human experts to ground truth their content and estimate error rates. All unchecked database entries will be flagged

as such, allowing users to decide how much to trust each data point that they extract, and the database user interface will allow for users to validate unchecked entries if and when they are able, facilitating the continuous increase in proportion of the database that is human validated. In parallel we will periodically implement semi-automated checks of the database contents using biological principles, either via automated scripts or again using AI. For instance, if a single ZW system is found in well-sampled clade that otherwise contains a conserved XY system with a single origin, it could be flagged for manual checking. This is one simple example of dozens of checks that could be implemented to reduce error rates in the database if needed.

The second challenge will be to integrate the information from the literature. This information will span multiple data types, a vast taxonomy, and come from sources spanning over a century. Encoding this information in a database in such a way that is consistent and usable will represent one of the largest challenges in the project. To achieve this, we are developing the Tree of Sex Ontology (TOSO <https://github.com/Tree-of-Sex/ToS-Ontology>), which will capture the hierarchy of terms within the database and their logical relationships. Using an ontology allows us to integrate information via a published and versioned logic, which will facilitate both data entry and interpretation. For example, consider a scenario in which a researcher wants to enter an observation of automixis for a given taxon. Automixis is a mode of parthenogenesis, thus the logic of TOSO will allow for automatic entry of “parthenogenesis” for reproductive strategy. Using an ontology also allows for logical testing of the consistency of the submitted data as well as identification of possible discrepancies between studies. Finally, the ontology has the capacity to record synonyms and give them context, facilitating the resolution of conflicting or overlapping naming conventions between research disciplines and taxonomic specialities. For example, in the plant literature “apomixis” refers to asexuality via unfertilized seeds, whereas, in the animal literature, “apomixis” refers to a type of asexuality without meiosis and without recombination (mixing) between haplotypes (mitotic asexuality) (Neiman et al., 2014; Van Dijk, 2009). The ontology allows contextual definitions of these terms that can be used in parallel, and users will be able to both submit data and query the database without enforcing any of their individual community’s standards on others. TOSO will therefore be the tool by which we integrate all information within the Tree of Sex and will help us unify this currently disparate field.

### Outlook

Compiling all knowledge relevant to the diversity of reproductive strategies in eukaryotes is ambitious but promises to advance the field of evolutionary biology. The integration of information across the taxa and topics will increase understanding of the patterning of reproductive strategies across the eukaryote phylogeny, identify gaps towards which we can direct future research, and facilitate large-scale comparative analyses to test fundamental evolutionary questions. However, this task will require a large community effort thus we warmly welcome researchers from all locations and backgrounds who are directly or indirectly interested in reproduction to join the Tree of Sex consortium ([treeofsex.org](http://treeofsex.org)).

## Glossary

Below are definitions of some important terms used in this article. Where possible we have adopted current definitions from existing biological ontologies (see term IDs). In some cases, we have slightly modified definitions to match our use of the term (see “Adapted from”). For terms lacking a definition in existing ontologies, or for which the existing definition(s) is inadequate, we have adopted new definitions based on our own current understanding.

We acknowledge that our interpretation of some of the terms below may reflect some of our biases as researchers,

and that some definitions may not include the full diversity of cases that exist in nature. Thus, the definitions below are presented to clarify our use of terms in this manuscript, and not as the “only” or “correct” definition for a given term.

We also note that definitions of terms may change from those presented below as our biological understanding expands. These changes will be reflected in the ontology generated by the consortium as part of the ongoing work that the Tree of Sex is conducting. This ontology, will include precise and inclusive definitions connected by logical relationships between terms, that will eventually be used for the database.

<b>Anisogamy</b>	A characteristic of sexually reproducing organisms whereby the different gamete types produced by a species differ substantially in size or form.
<b>Arrhenotoky</b>	A sex determination strategy in which males develop from unfertilized eggs and are haploid, and females develop from fertilized eggs and are diploid. From <a href="#">GSSO:000121</a>
<b>Asexual reproduction</b>	A type of reproduction in which new individuals are produced from a single organism, either from an unfertilized egg or from a single cell or group of cells. From <a href="#">GO:0019954</a>
<b>Automixis</b>	In the context of fungal reproduction, the fusing gametes might be from the same meiotic tetrad. In plants and animals, it refers to a form of parthenogenesis, where the progeny is not genetically identical to the parent.
<b>Dioecy</b> (synonym: <b>Gonochorism</b> )	Sexual strategy in which anisogamous (macro or micro) gametes are produced in separate individuals. Dioecy is used in plants while gonochorism typically refers to animals.
<b>Dioicy</b>	When sex is determined in the haploid stage, the occurrence of gametophytes that produce only one type of gametes; separate sexes.
<b>Environmental sex determination (ESD)</b>	The determination of sex by a non-genetic cue, such as temperature, nutrient availability, or social context. Adapted from <a href="#">GSSO:000107</a>
<b>Female</b>	In anisogamous organisms, a category of individual or organ (e.g., flower) sex which produces macrogametes (ova). Note that this simplified categorization may not be reflective of existing phenotypic diversity as it ignores the complex and often overlapping combination of other traits associated with sex.
<b>Gametes</b>	A mature sexual reproductive cell often (but not always) having a single set of chromosomes (haploid). Adapted from <a href="#">CL:0000300</a>
<b>Gametolog</b>	One of a homologous set of genes in the non-recombining region on the sex chromosomes. Can also be used to refer to the individual sex chromosomes themselves (e.g., the X and Y chromosomes are the two gametologs of an XY system).
<b>Gametophyte</b>	The multicellular haploid stage of the life cycle of a (sexual) plant species in which gametes are produced.
<b>Genotypic sex determination (GSD)</b>	Determination of sex by genotypic constitution, namely by chromosomal or genetic factors consistently associated with each sex. Adapted from <a href="#">GSSO:000104</a>
<b>Gonochorism</b> (synonym: <b>Dioecy</b> )	Sexual strategy in which anisogamous (macro or micro) gametes are produced in separate individuals. Gonochorism typically refers to animals, whereas dioecy is used in plants.
<b>Haploid</b>	Chromosomally reduced stage of the life cycle. In some species, the reduced stage is the relatively short phase with gametes directly produced during meiosis; in others (fungi, mosses, ferns) meiosis produce sporophytes that undergo mitotic divisions to form a multicellular body before gametes are produced. Note that haploid is often used to specifically refer to a cell containing a single set of chromosomes (i.e., monoploid), though here we use it only to mean reduced.
<b>Hermaphroditism</b>	An umbrella term for a suite of sexual reproductive strategies whereby individuals can produce both macro and microgametes. In plants, hermaphroditism refers specifically to plants with flowers that can produce both male and female gametes, and is distinct from Monoecy (see below). See also “Sequential hermaphroditism” and “Simultaneous hermaphroditism.”
<b>Heterogametic</b>	A characteristic of an individual producing gametes that differ in sex chromosome constitution, such as heterogametic males in XX/XY (or females in ZZ/ZW systems) that produce X- or Y-containing sperm (or Z- or W-containing eggs).
<b>Heteromorphic sex chromosomes</b>	A sex chromosome system in which the sex chromosomes (e.g., X and Y) are morphologically different from each other when observed via microscopy.
<b>Homogametic</b>	A characteristic of an individual producing gametes with identical sex chromosome constitution, such as females in systems (XX/XY) (or males in ZZ/ZW systems), that produce exclusively X-containing eggs (or Z-containing sperm).
<b>Homomorphic sex chromosomes</b>	A sex chromosome system in which the sex chromosomes (e.g., X and Y) are morphologically indistinguishable when observed via microscopy.
<b>Intragametophytic selfing</b>	A form of self-fertilization between gametes produced within the same gametophyte.
<b>Isogamy</b>	A characteristic of sexually reproducing organisms whereby the different gamete types produced by a species do not differ substantially in size or shape.
<b>Male</b>	In the context of anisogamy, a category of individual or organ (e.g., flower) sex that produces microgametes (sperm). Note that this simplified categorization may not be reflective of existing phenotypic diversity as it ignores the complex and often overlapping combination of other traits associated with sex.
<b>Mating type</b>	A characteristic of gametes or individuals of sexually reproducing (often isogamous) species that controls their compatibility for genetic mixing (fusion or mating, respectively).

<b>Monoecy</b>	A reproductive strategy In plants, wherein individual plants possess both male and female flowers. Note that this is distinct from “hermaphroditism.”
<b>Monoicy</b>	When sex is determined in the haploid stage, the occurrence of gametophytes that produce both male and female gametes.
<b>Neo-sex chromosome</b>	A sex chromosome resulting from the fusion of either gametolog of an existing sex chromosome pair (e.g., X or Y to an autosome, which generates a Neo-X chromosome, or a Neo-Y chromosome).
<b>Parthenogenesis</b>	Development of an egg into an embryo without being fertilized. <a href="#">MP:0009443</a>
<b>Programmed DNA elimination</b>	A process in which genomic fragments or entire chromosomes are eliminated from somatic cells. <a href="#">GO:0031049</a>
<b>Pseudoautosomal region</b>	The region(s) of a sex chromosome which still recombine(s).
<b>Reproductive strategy</b>	The set of traits (morphological, physiological, behavioral, etc.), that help an organism reproduce.
<b>Self-compatibility</b>	The opposite of “Self incompatibility,” i.e., gametes of different types from the same organism are able to fuse, resulting in reproduction.
<b>Self-incompatibility</b>	A self-recognition mechanism reducing self-fertilization and inbreeding. Adapted from <a href="#">TO:0000310</a>
<b>Sequential hermaphroditism</b>	Sexual reproductive system of some taxa whereby individuals switch between producing either macro or microgametes at some point in their lifetime
<b>Sex chromosome</b>	A chromosome harbouring a locus involved in sex determination. Sex linkage within a sex chromosome may be restricted to this locus, or extend, via recombination suppression, across varying proportions of its length. Adapted from <a href="#">GO:0000803</a> .
<b>Sex</b>	A category of individual, tissue, genotype, or gamete in anisogamous taxa that undergo sexual reproduction. Applied based on primary or secondary sexual characteristics. Common categories for sexes include female and male, although these generally represent a simplification of phenotypic diversity and continuous variation. Adapted from <a href="#">PATO:0000047</a> . Sometimes akin to mating type in isogamous organisms.
<b>Sexual reproduction</b>	A type of reproduction in anisogamous or isogamous eukaryotes that combines the genetic material of two gametes, from two individuals or from a single individual. Adapted from <a href="#">GO:0019953</a> .
<b>Sexual system</b>	The pattern of sex allocation and/or mating behaviour across individuals in a sexually reproducing species (e.g., gonochorism, hermaphroditism, mixed sexual systems). Adapted from <a href="#">GSSO:011864</a>
<b>Simultaneous hermaphrodites</b>	Sexual reproductive system of some taxa whereby an individual is capable of producing both macro and microgametes at the same time.
<b>Syngamy</b>	The union of gametes of types to form a zygote during sexual reproduction. Syngamy involves the fusion of cytoplasm (plasmogamy) and nuclei (karyogamy). Some species live a substantial proportion of their lives in between these two stages (fungi). From <a href="#">GO:0009566</a>

## Data availability

No new data are presented in this manuscript. Details of the analysis in Figure 1 can be found in the legend.

## Author contributions

Daniel Jeffries (Conceptualization [lead], Formal analysis [lead], Visualization [lead], Writing—original draft [lead], Writing—review & editing [lead]), Chiara Benvenuto (Writing—review & editing [supporting]), Astrid Böhne (Writing—review & editing [supporting]), Christelle Fraïsse (Formal analysis [lead], Writing—original draft [equal], Writing—review & editing [equal]), Sònia Garcia (Writing—original draft [equal], Writing—review & editing [equal]), Paul Jay (Writing—original draft [equal], Writing—review & editing [equal]), Lukáš Kratochvíl (Writing—original draft [equal], Writing—review & editing [equal]), Caitlin E. McDonough-Goldstein (Writing—review & editing [supporting]), Aurora Ruiz-Herrera (Writing—original draft [equal], Writing—review & editing [equal]), Cibele G. Sotero-Caio (Conceptualization [supporting], Visualization [supporting], Writing—original draft [equal], Writing—review & editing [equal]), Nicole Valenzuela (Writing—original draft [equal], Writing—review & editing [equal]), Melissa Wilson (Writing—original draft [equal], Writing—review & editing [equal]), and Kamil S. Jaron (Conceptualization [supporting],

Formal analysis [lead], Visualization [lead], Writing—original draft [lead], Writing—review & editing [lead])

## Funding

C.G.S.C. and K.S.J. were supported by the Wellcome Trust grant number 220540. L.K. is supported by the Czech Science Foundation (23-07665S). A.R.H. is supported by the Spanish Ministry of Science and Innovation (PID2020-112557GB-I00 funded by AEI/10.13039/501100011033), the Agència de Gestió d'Ajuts Universitaris i de Recerca, AGAUR (2021SGR00122) and the Catalan Institution for Research and Advanced Studies (ICREA). P.J. is supported by a Human Frontier Science Program (HFSP) fellowship (LT0033/2022-L). C.E.M.G. is supported by a National Science Foundation postdoctoral research fellowship (DBI 2208973). D.L.J. is supported by the Swiss National Science Foundation project grant TMAG-3\_209309. A.B. is supported by the Deutsche Forschungsgemeinschaft DFG grant IDs 492407022 and 497674620 / 1. C.B. could attend the Tree of Sex workshops thanks to an ASAB (Association for the Study of Animal Behaviour) grant. M.A.W. is supported by the NIGMS of the National Institutes of Health award R35GM124827 to M.A.W. S.G. is supported by the Spanish Ministry of Science and Innovation (CNS2023-143604 funded by AEI/10.13039/501100011033) and the Agència de Gestió d'Ajuts Universitaris i de Recerca, AGAUR (2021SGR00315).

N.V. is supported in part by grant IOS 2127995 from the USA National Science Foundation.

## Acknowledgments

We would like to acknowledge the support of Enabling Platforms, Tree of Life Programme, Wellcome Sanger Institute in providing the Tree of Sex infrastructure such as the database, its interface and our webpage. We also thanks the editors and reviewers for for their many useful comments.

## Conflicts of interest

None declared.

## Tree of Sex Consortium:

Aleksandra Bliznina, Valentina Peona, Thomas Desvignes, Aparna Lajmi, Christina N. Hodson, Yann Guiguen, Mónica Moura, Paul Jay, Tanja Slotte, Bjarte H. Jordal, Michael A. White, Philipp Schiffer, Maja Lazarević, Christopher G. Wilson, Joana Isabel Meier, Joris M. Koene, Christelle Fraisse, Mathias Scharmann, Bahadır Onsoy, Jun Kitano, Giulia Lin, Ann Kathrin Huylmans, Emmarie P. Alexander, Caitlin E. McDonough-Goldstein, Sara Calhim, Ashwini V. Mohan, Shana Pau, Sophie Helen Smith, Cecile Molinier, Melissah Rowe, Leo W. Beukeboom, Stephen I. Wright, Matjaž Kuntner, John R. Pannell, Ludovic Dutoit, Edina Nemesházi, Tony Gamble, Peter Szoenyi, Petra Bulankova, Vratislav Peska, Caleb J. Krueger, Astrid Böhne, Dragana Cvetković, Mércia Patrícia Pereira Silva, Thaís Elias Almeida, Frédéric Veyrunes, Agnieszka Lipinska, Veronika Bókony, Jana C. Vamosi, Peter Glasnović, J. Antonio Baeza, Melissa A. Wilson, Tatiana Giraud, Cibele G. Sotero-Caio, Stuart V. Nielsen, Sònia Garcia, Richard Challis, Lukáš Kratochvíl, Luohao Xu, Marta Turon, Jie Wang, Deborah Charlesworth, Charlotte J. Wright, Bozena Kolano, Brendan J. Pinto, Giacomo Potente, Wagner Luiz dos Santos, Zoran Marčić, Clare E. Holleley, Soleil E. Young, Stuart F. McDaniel, Lucija Andjel, Matthew W. Hahn, Claudia C. Weber, Fredric J. Janzen, Lewis Stevens, Ahammad Kabir, Jennifer D. Gresham, Paul A. Saunders, Marine Arakelyan, Jiří Král, Duminda S.B. Dissanayake, Stacy A. Krueger-Hadfield, Qi Zhou, Jonathan M.D. Wood, Catherine L. Peichel, Karel Janko, Maria Immacolata Ferrante, Chiara Benvenuto, Michail Rovatsos, Andrew J. Mongue, Máire Ní Leathlobhair, Jessica K. Abbott, Jonna Kulmuni, Anna Torgasheva, Alexander Suh, David H. Lunt, Susana M. Coelho, Sam Ebdon, Alex Makunin, Melissa A. Touns, Jens Bast, Quentin Helleu, Quentin Helleu, Tony Heitkam, Vladimir Trifonov, Sadye Paez, Rainer Melzer, Natalia Borowska-Zuchowska, Nicole Valenzuela, Daniela H. Palmer Droguett, Rita Monteiro, Petr Nguyen, Kamil S. Jaron, Daniel L. Jeffries, Caroline Howard, Tanja Schwander, Kerstin Howe, Ravinder K. Kanda, Arthur Georges, Wen-Juan Ma, Ore Francis, Mark Blaxter, Sarah P. Otto, Roman Hobza, Louise D. Heitzmann, Vasiliki Kousteni, Aurora Ruiz-Herrera, David Peris, James Umen, Cristiana Ramalho Maciel, Alexandr Sember, Ricardo Utsunomia, Yukako Katsura, Matthias Stöck, Simon Kershenbaum, Tymoteusz Pieszkowski, Vilma Loreto, Martha Mercy Mulongo. All contributors listed above gave their consent to be cited in the collaborative group of the Tree of Sex Consortium.

## References

- Aanen, D., Beekman, M., & Kokko, H. (2016). Weird sex: The underappreciated diversity of sexual reproduction. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 371(1706), 20160262. <https://doi.org/10.1098/rstb.2016.0262>
- Aghi, K., Anderson, B. M., Castellano, B. M., ... Weinberg, Z. Y. (2024). Rigorous science demands support of transgender scientists. *Cell*, 187(6), 1327–1334. <https://doi.org/10.1016/j.cell.2024.02.021>
- Ahmed, S., Cock, J. M., Pessia, E., ... Coelho, S. M. (2014). A haploid system of sex determination in the brown alga *Ectocarpus* sp. *Current Biology: CB*, 24(17), 1945–1957.
- Ahnesjö, I., Brealey, J. C., Günter, K. P., & ... Vasconcelos, P. (2020). Considering gender-biased assumptions in evolutionary biology. *Evolutionary Biology*, 47(1), 1–5. <https://doi.org/10.1007/s11692-020-09492-z>
- Alonzo, S. H. (2010). Social and coevolutionary feedbacks between mating and parental investment. *Trends in Ecology & Evolution*, 25(2), 99–108. <https://doi.org/10.1016/j.tree.2009.07.012>
- Álvarez-González, L., Burden, F., Doddamani, D., ... Ruiz-Herrera, A. (2022). 3D chromatin remodelling in the germ line modulates genome evolutionary plasticity. *Nature Communications*, 13(1), 2608. <https://doi.org/10.1038/s41467-022-30296-6>
- Álvarez-González, L., & Ruiz-Herrera, A. (2025). Evolution of 3D chromatin folding. *Annual Review of Animal Biosciences*, 13(1), 49–71. <https://doi.org/10.1146/annurev-animal-111523-102233>
- Anderson, N. W., Hjelmén, C. E., & Blackmon, H. (2020). The probability of fusions joining sex chromosomes and autosomes. *Biology Letters*, 16(11), 20200648. <https://doi.org/10.1098/rsbl.2020.0648>
- Ansai, S., Montenegro, J., Masengi, K. W. A., ... Kitano, J. (2022). Diversity of sex chromosomes in Sulawesi medaka fishes. *Journal of Evolutionary Biology*, 35(12), 1751–1764. <https://doi.org/10.1111/jeb.14076>
- Araujo, D., Schneider, M. C., Paula-Neto, E., & Cella, D. (2012). Sex chromosomes and meiosis in spiders: A review. In A. Swan (Ed.), *Meiosis—Molecular mechanisms and cytogenetic diversity* (pp. 87–108). InTechOpen.
- Araujo, D., Schneider, M. C., Paula-Neto, E., & Cella, D. M. (2024). The spider cytogenetic database. [www.arthropodacytogenetics.bio.br/spiderdatabase](http://www.arthropodacytogenetics.bio.br/spiderdatabase)
- Bachtrog, D., Kirkpatrick, M., Mank, J. E., ... Valenzuela, N. (2011). Are all sex chromosomes created equal? *Trends in Genetics: TIG*, 27(9), 350–357. <https://doi.org/10.1016/j.tig.2011.05.005>
- Bachtrog, D., Mahajan, S., & Bracewell, R. (2019). Massive gene amplification on a recently formed *Drosophila* Y chromosome. *Nature Ecology & Evolution*, 3(11), 1587–1597. <https://doi.org/10.1038/s41559-019-1009-9>
- Bachtrog, D., Mank, J. E., Peichel, C. L., ... Vamosi, J. C.; Tree of Sex Consortium (2014). Sex determination: Why so many ways of doing it? *PLoS Biology*, 12(7), e1001899. <https://doi.org/10.1371/journal.pbio.1001899>
- Baker, H. G. (1955). Self-compatibility and establishment after “long-distance” dispersal. *Evolution; International Journal of Organic Evolution*, 9(3), 347–349.
- Baránková, S., Pascual-Díaz, J. P., Sultana, N., ... Garcia, S. (2020). Sex-chrom, a database on plant sex chromosomes. *The New Phytologist*, 227(6), 1594–1604. <https://doi.org/10.1111/nph.16635>
- Baroiller, J. F., Chourrout, D., Fostier, A., & Jalabert, B. (1995). Temperature and sex chromosomes govern sex ratios of the mouthbrooding cichlid fish *Oreochromis niloticus*. *Journal of Experimental Zoology*, 273(3), 216–223. <https://doi.org/10.1002/jez.1402730306>
- Barrera-Redondo, J., Lipinska, A. P., Liu, P., ... Coelho, S. M. (2024). Origin and evolutionary trajectories of brown algal sex chromosomes. *bioRxiv* 575685. <https://doi.org/10.1101/2024.01.15.575685>, January 16, 2024, preprint: not peer reviewed.
- Barrett, S. C. H. (1998). The evolution of mating strategies in flowering plants. *Trends in Plant Science*, 3(9), 335–341.
- Bell, G. (1981). *The masterpiece of nature: Evolution and genetics of sexuality*. Kluwer Academic.

- Benvenuto, C., & Weeks, S. C. (2020). Hermaphroditism and gonochorism. In R. Cothran, & M. Thiel (Eds.), *The natural history of the Crustacea, Vol. VI: Reproductive biology* (pp. 197–241). Oxford University Press.
- Bergero, R., & Charlesworth, D. (2019). Reply to Wright *et al.*: How to explain the absence of extensive Y-specific regions in the guppy sex chromosomes. *Proceedings of the National Academy of Sciences of the United States of America*, 116(26), 12609–12610. <https://doi.org/10.1073/pnas.1906633116>
- Bergero, R., Gardner, J., Bader, B., ... Charlesworth, D. (2019). Exaggerated heterochiasmy in a fish with sex-linked male coloration polymorphisms. *Proceedings of the National Academy of Sciences of the United States of America*, 116(14), 6924–6931. <https://doi.org/10.1073/pnas.1818486116>
- Bernstein, H., & Bernstein, C. (2010). Evolutionary origin of recombination during meiosis. *Bioscience*, 60(7), 498–505. <https://doi.org/10.1525/bio.2010.60.7.5>
- Berzins, I. K., Yanong, R. P. E., LaDouceur, E. E. B., & Peters, E. C. (2021). Cnidaria. In LaDouceur, E. E. B. (Ed), *Invertebrate histology* (pp. 55–86). Wiley. <https://doi.org/10.1002/9781119507697.ch3>
- Beukeboom, L. W., & Perrin, N. (2014). *The evolution of sex determination*. Oxford University Press.
- Beukeboom, L. W., & Vrijenhoek, R. C. (1998). Evolutionary genetics and ecology of sperm-dependent parthenogenesis. *Journal of Evolutionary Biology*, 11(6), 755–782. <https://doi.org/10.1007/s000360050117>
- Beye, M., Hasselmann, M., Fondrk, M. K., ... Omholt, S. W. (2003). The gene *csd* is the primary signal for sexual development in the honeybee and encodes an SR-type protein. *Cell*, 114(4), 419–429. [https://doi.org/10.1016/S0092-8674\(03\)00606-8](https://doi.org/10.1016/S0092-8674(03)00606-8)
- Billiard, S., López-Villavicencio, M., Devier, B., ... Giraud, T. (2011). Having sex, yes, but with whom? Inferences from fungi on the evolution of anisogamy and mating types. *Biological Reviews of the Cambridge Philosophical Society*, 86(2), 421–442. <https://doi.org/10.1111/j.1469-185X.2010.00153.x>
- Birky, C. W. Jr (1996). Heterozygosity, heteromorphy, and phylogenetic trees in asexual eukaryotes. *Genetics*, 144(1), 427–437. <https://doi.org/10.1093/genetics/144.1.427>
- Bista, B., González-Rodelas, L., Álvarez-González, L., ... Valenzuela, N. (2024). De novo genome assemblies of two cryptodiran turtles with ZZ/ZW and XX/XY sex chromosomes provide insights into patterns of genome reshuffling and uncover novel 3D genome folding in amniotes. *Genome Research*, 34(10), 1553–1569. <https://doi.org/10.1101/gr.279443.124>
- Blackmon, H., & Demuth, J. P. (2014). Estimating tempo and mode of Y chromosome turnover: Explaining Y chromosome loss with the fragile Y hypothesis. *Genetics*, 197(2), 561–572. <https://doi.org/10.1534/genetics.114.164269>
- Blackmon, H., & Demuth, J. P. (2015a). Coleoptera karyotype database. *The Coleopterists Bulletin*, 69(1), 174–175. <https://doi.org/10.1649/0010-065x-69.1.174>
- Blackmon, H., & Demuth, J. P. (2015b). The fragile Y hypothesis: Y chromosome aneuploidy as a selective pressure in sex chromosome and meiotic mechanism evolution. *BioEssays*, 37(9), 942–950. <https://doi.org/10.1002/bies.201500040>
- Blackmon, H., Hardy, N. B., & Ross, L. (2015). The evolutionary dynamics of haplodiploidy: Genome architecture and haploid viability. *Evolution; International Journal of Organic Evolution*, 69(11), 2971–2978. <https://doi.org/10.1111/evo.12792>
- Blackmon, H., Justison, J., Mayrose, I., & Goldberg, E. E. (2019). Meiotic drive shapes rates of karyotype evolution in mammals. *Evolution; International Journal of Organic Evolution*, 73(3), 511–523. <https://doi.org/10.1111/evo.13682>
- Blackmon, H., Ross, L., & Bachtrog, D. (2017). Sex determination, sex chromosomes, and karyotype evolution in insects. *The Journal of Heredity*, 108(1), 78–93. <https://doi.org/10.1093/jhered/esw047>
- Blaser, O., Neuenschwander, S., & Perrin, N. (2014). Sex-chromosome turnovers: The hot-potato model. *The American Naturalist*, 183(1), 140–146. <https://doi.org/10.1086/674026>
- Blaxter, M., Pauperio, J., Schoch, C., & Howe, K. (2024b). Taxonomy Identifiers (TaxId) for Biodiversity Genomics: A guide to getting TaxId for submission of data to public databases. *Wellcome Open Research*, 9, 591. <https://doi.org/10.12688/wellcomeopenres.22949.1>
- Blaxter, M. L., Leech, C., & Lunt, D. H. (2024a). A catalogue of chromosome counts for Phylum Nematoda. <https://wellcomeopenresearch.org/articles/9-55>
- Bókony, V., Kövér, S., Nemesházi, E., ... Székely, T. (2017). Climate-driven shifts in adult sex ratios via sex reversals: The type of sex determination matters. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 372(1729), 20160325. <https://doi.org/10.1098/rstb.2016.0325>
- Bornmann, L., Haunschild, R., & Mutz, R. (2021). Growth rates of modern science: A latent piecewise growth curve approach to model publication numbers from established and new literature databases. *Humanities and Social Sciences Communications*, 8(1), 1–15.
- Bracewell, R., Tran, A., Chatla, K., & Bachtrog, D. (2024). Sex and neo-sex chromosome evolution in beetles. *PLoS Genetics*, 20(11), e1011477. <https://doi.org/10.1371/journal.pgen.1011477>
- Brandt, A., Tran Van, P., Bluhm, C., ... Bast, J. (2021). Haplotype divergence supports long-term asexuality in the oribatid mite *Oppiella nova*. *Proceedings of the National Academy of Sciences of the United States of America*, 118(38), e2101485118. <https://doi.org/10.1073/pnas.2101485118>
- Breton, S., Capt, C., Guerra, D., & Stewart, D. (2018). Sex-determining mechanisms in bivalves. In Leonard, J. (Ed), *Transitions between sexual systems* (pp. 165–192). Springer International Publishing.
- Bridges, C. B. (1916). Non-disjunction as proof of the chromosome theory of heredity. *Genetics*, 1(1), 1–52. <https://doi.org/10.1093/genetics/1.1.1>
- Brush, S. G. (1978). Nettie M. Stevens and the discovery of sex determination by chromosomes. *Isis; an International Review Devoted to the History of Science and Its Cultural Influences*, 69(247), 163–172. <https://doi.org/10.1086/352001>
- Bull, J. J. (1978). Sex-chromosomes in haploid dioecy; unique contrast to Muller's theory for diploid dioecy. *American Naturalist*, 112(983), 245–250. <https://doi.org/10.1086/283267>
- Bull, J. J. (1983). *Evolution of sex determining mechanisms*. The Benjamin/Cummings Publishing Company, Inc.
- Carey, S. B., Jenkins, J., Lovell, J. T., ... McDaniel, S. F. (2021). Genrich UV sex chromosomes harbor conserved regulators of sexual development. *Science Advances*, 7(27), eabh2488. <https://doi.org/10.1126/sciadv.abh2488>
- Carvalho, A. B., Dobo, B. A., Vibrantovski, M. D., & Clark, A. G. (2001). Identification of five new genes on the Y chromosome of *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences of the United States of America*, 98(23), 13225–13230. <https://doi.org/10.1073/pnas.231484998>
- Cerepaka, C., & Schlupp, I. (2023). Sperm specificity and potential paternal effects in gynogenesis in the Amazon Molly (*Poecilia formosa*). *PeerJ*, 11(e16118), e16118. <https://doi.org/10.7717/peerj.16118>
- Challis, R., Kumar, S., Sotero-Caio, C., ... Blaxter, M. (2023). Genomes on a Tree (GoaT): A versatile, scalable search engine for genomic and sequencing project metadata across the eukaryotic tree of life. *Wellcome Open Research*, 8, 24. <https://doi.org/10.12688/wellcomeopenres.18658.1>
- Chalopin, D., Volff, J. -N., Galiana, D., ... Scharl, M. (2015). Transposable elements and early evolution of sex chromosomes in fish. *Chromosome Research*, 23(3), 545–560. <https://doi.org/10.1007/s10577-015-9490-8>
- Charlesworth, B. (1978). The population genetics of anisogamy. *Journal of Theoretical Biology*, 73(2), 347–357. [https://doi.org/10.1016/0022-5193\(78\)90195-9](https://doi.org/10.1016/0022-5193(78)90195-9)
- Charlesworth, B., Coyne, J. A., & Barton, N. H. (1987). The relative rates of evolution of sex chromosomes and autosomes. *The American Naturalist*, 130(1), 113–146. <https://doi.org/10.1086/284701>

- Charlesworth, B., & Olito, C. (2024). Making sense of recent models of the “sheltering” hypothesis for recombination arrest between sex chromosomes. *Evolution; International Journal of Organic Evolution*, 78(12), 1891–1899. <https://doi.org/10.1093/evolut/qpae119>
- Charlesworth, B., Sniegowski, P., & Stephan, W. (1994). The evolutionary dynamics of repetitive DNA in eukaryotes. *Nature*, 371(6494), 215–220. <https://doi.org/10.1038/371215a0>
- Charlesworth, D. (2022). The mysterious sex chromosomes of haploid plants. *Heredity*, 129(1), 17–21. <https://doi.org/10.1038/s41437-022-00524-2>
- Charlesworth, D. (2023). Why and how do Y chromosome stop recombining? *Journal of Evolutionary Biology*, 36(3), 632–636. <https://doi.org/10.1111/jeb.14137>
- Charlesworth, D., Bergero, R., Graham, C., ... Yong, L. (2020). Locating the sex determining region of linkage group 12 of guppy (*Poecilia reticulata*). *G3 (Bethesda, Md.)*, 10(10), 3639–3649. <https://doi.org/10.1534/g3.120.401573>
- Charlesworth, D., & Charlesworth, B. (1980). Sex differences in fitness and selection for centric fusions between sex-chromosomes and autosomes. *Genetical Research*, 35(2), 205–214. <https://doi.org/10.1017/s0016672300014051>
- Charlesworth, D., Charlesworth, B., & Marais, G. (2005). Steps in the evolution of heteromorphic sex chromosomes. *Heredity*, 95(2), 118–128. <https://doi.org/10.1038/sj.hdy.6800697>
- Chen, M. -S., Niu, L., Zhao, M. -L., ... Xu, Z. -F. (2020). De novo genome assembly and Hi-C analysis reveal an association between chromatin architecture alterations and sex differentiation in the woody plant *Jatropha curcas*. *GigaScience*, 9(2), gaa009. <https://doi.org/10.1093/gigascience/giaa009>
- Chen, X. I., Mei, Y., Chen, M., ... Li, F. (2021). InSexBase: An annotated genomic resource of sex chromosomes and sex-biased genes in insects. *Database: The Journal of Biological Databases and Curation*, 2021, baab001. <https://doi.org/10.1093/database/baab001>
- Cheptou, P. -O. (2018). Does the evolution of self-fertilization rescue populations or increase the risk of extinction? *Annals of Botany*, 123(2), 337–345. <https://doi.org/10.1093/aob/mcy144>
- Chepurnov, V. A., Mann, D. G., Sabbe, K., & Vyverman, W. (2004). Experimental studies on sexual reproduction in diatoms. *International Review of Cytology*, 237, 91–154. [https://doi.org/10.1016/S0074-7696\(04\)37003-8](https://doi.org/10.1016/S0074-7696(04)37003-8)
- Choleva, L., Apostolou, A., Rab, P., & Janko, K. (2008). Making it on their own: Sperm-dependent hybrid fishes (*Cobitis*) switch the sexual hosts and expand beyond the ranges of their original sperm donors. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 363(1505), 2911–2919. <https://doi.org/10.1098/rstb.2008.0059>
- Choleva, L., & Janko, K. (2013). Rise and persistence of animal polyploidy: Evolutionary constraints and potential. *Cytogenetic and Genome Research*, 140(2-4), 151–170. <https://doi.org/10.1159/000353464>
- Church, S. H., Donoughe, S., de Medeiros, B. A. S., & Extavour, C. G. (2019). A dataset of egg size and shape from more than 6,700 insect species. *Scientific Data*, 6(1), 104. <https://doi.org/10.1038/s41597-019-0049-y>
- Coelho, S. M., Gueno, J., Lipinska, A. P., ... Umen, J. G. (2018). UV chromosomes and haploid sexual systems. *Trends in Plant Science*, 23(9), 794–807. <https://doi.org/10.1016/j.tplants.2018.06.005>
- Coffing, G. C., Tittes, S., Small, S. T., ... Kern, A. D. (2024). Cephalopod sex determination and its ancient evolutionary origin revealed by chromosome-level assembly of the California two-spot octopus. *bioRxiv* 581452, <https://doi.org/10.1101/2024.02.21.581452>, February 21, 2024, preprint: not peer reviewed.
- Colaco, S., & Modi, D. (2018). Genetics of the human Y chromosome and its association with male infertility. *Reproductive Biology and Endocrinology: RB&E*, 16(1), 14. <https://doi.org/10.1186/s12958-018-0330-5>
- Collin, R. (2006). Sex ratio, life-history invariants, and patterns of sex change in a family of protandrous gastropods. *Evolution; International Journal of Organic Evolution*, 60(4), 735–745.
- Constable, G. W. A., & Kokko, H. (2021). Parthenogenesis and the evolution of anisogamy. *Cells*, 10(9), 2467. <https://doi.org/10.3390/cells10092467>
- Cordellier, M., Schneider, J. M., Uhl, G., & Posnien, N. (2020). Sex differences in spiders: from phenotype to genomics. *Development Genes and Evolution*, 230(2), 155–172. <https://doi.org/10.1007/s00427-020-00657-6>
- Cossard, G. G., & Pannell, J. R. (2019). A functional decomposition of sex inconstancy in the dioecious, colonizing plant *Mercurialis annua*. *American Journal of Botany*, 106(5), 722–732. <https://doi.org/10.1002/ajb2.1277>
- Costa, W. J. E. M., Lima, S. M. Q., & Bartolette, R. (2010). Androdioecy in *Kryptolebias* killifish and the evolution of self-fertilizing hermaphroditism. *Biological Journal of the Linnean Society*, 99(2), 344–349. <https://doi.org/10.1111/j.1095-8312.2009.01359.x>
- Coyne, J. A. (1985). The genetic basis of Haldane’s rule. *Nature*, 314(6013), 736–738. <https://doi.org/10.1038/314736a0>
- Coyne, J. A. (2018). “two rules of speciation” revisited. *Molecular Ecology*, 27(19), 3749–3752. <https://doi.org/10.1111/mec.14790>
- Crossman, A., & Charlesworth, D. (2014). Breakdown of dioecy: Models where males acquire coxexual functions. *Evolution; International Journal of Organic Evolution*, 68(2), 426–440. <https://doi.org/10.1111/evo.12283>
- D’Ambrosio, U., Alonso-Lifante, M., Barros, K., ... Garcia, S. (2017). B-chrom: A database on B-chromosomes of plants, animals and fungi. *The New Phytologist*, 216(3), 635–642.
- da Silva, J. (2022). Gene dynamics of haplodiploidy favor eusociality in the Hymenoptera. *Evolution; International Journal of Organic Evolution*, 76(7), 1546–1555. <https://doi.org/10.1111/evo.14518>
- Dapper, A. L., & Payseur, B. A. (2017). Connecting theory and data to understand recombination rate evolution. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 372(1736), 20160469. <https://doi.org/10.1098/rstb.2016.0469>
- de la Filia, A. G., Bain, S. A., & Ross, L. (2015). Haplodiploidy and the reproductive ecology of Arthropods. *Current Opinion in Insect Science*, 9, 36–43. <https://doi.org/10.1016/j.cois.2015.04.018>
- de Vries, C., & Lehtonen, J. (2023). Sex-specific assumptions and their importance in models of sexual selection. *Trends in Ecology & Evolution*, 38(10), 927–935. <https://doi.org/10.1016/j.tree.2023.04.013>
- Deakin, J. E., Potter, S., O’Neill, R., Kratochvíl, L., ... Ezaz, T. (2019). Chromosomics: Bridging the gap between genomes and chromosomes. *Genes*, 10(8), 627. <https://doi.org/10.3390/genes10080627>
- Debus, B. (1978). “Nodules” in the achiasmatic meiosis of *Bithynia* (Mollusca, Prosobranchia). *Chromosoma*, 69(1), 81–92. <https://doi.org/10.1007/BF00327382>
- Dedukh, D., Altmanová, M., Petrosyan, R., ... Kratochvíl, L. (2024). Premeiotic endoreplication is the mechanism of obligate parthenogenesis in rock lizards of the genus *Darevskia*. *Biology Letters*, 20(9), 20240182. <https://doi.org/10.1098/rsbl.2024.0182>
- Dedukh, D., da Cruz, I., Kneitz, S., ... Scharl, M. (2022). Achiasmatic meiosis in the unisexual Amazon molly, *Poecilia formosa*. *Chromosome Research*, 30(4), 443–457. <https://doi.org/10.1007/s10577-022-09708-2>
- Deveson, I. W., Holleley, C. E., Blackburn, J., ... Georges, A. (2017). Differential intron retention in Jumonji chromatin modifier genes is implicated in reptile temperature-dependent sex determination. *Science Advances*, 3(6), e1700731. <https://doi.org/10.1126/sciadv.1700731>
- Dolmatov, I. Y., Afanasyev, S. V., & Boyko, A. V. (2018). Molecular mechanisms of fission in echinoderms: Transcriptome analysis. *PLoS One*, 13(4), e0195836. <https://doi.org/10.1371/journal.pone.0195836>
- Duffy, T. A., Hice, L. A., & Conover, D. O. (2015). Pattern and scale of geographic variation in environmental sex determination in the Atlantic silverside, *Menidia menidia*. *Evolution; International Journal of Organic Evolution*, 69(8), 2187–2195. <https://doi.org/10.1111/evo.12724>

- Dyer, A. F. (1979). The experimental biology of ferns. *Transactions of the Botanical Society of Edinburgh*, 43(2), 75–90. <https://doi.org/10.1080/03746607908685341>
- Ehlers, B. K., & Bataillon, T. (2007). “Inconstant males” and the maintenance of labile sex expression in subdioecious plants. *The New Phytologist*, 174(1), 194–211. <https://doi.org/10.1111/j.1469-8137.2007.01975.x>
- El Taher, A., Ronco, F., Matschiner, M., ... Böhne, A. (2021). Dynamics of sex chromosome evolution in a rapid radiation of cichlid fishes. *Science Advances*, 7(36), eabe8215. <https://doi.org/10.1126/sciadv.abe8215>
- Engelstädter, J. (2017). Asexual but not clonal: Evolutionary processes in autotictic populations. *Genetics*, 206(2), 993–1009. <https://doi.org/10.1534/genetics.116.196873>
- Ewers-Saucedo, C., Hope, N. B., & Wares, J. P. (2016). The unexpected mating system of the androdioecious barnacle *Chelonibia testudinaria* (Linnaeus 1758). *Molecular Ecology*, 25(9), 2081–2092. <https://doi.org/10.1111/mec.13593>
- Fausto-Sterling, A. (2008). *Sexing the body: Gender politics and the construction of sexuality*. Basic Books.
- Feldmeyer, B., Kozielska, M., Kuijper, B., ... Pen, I. (2008). Climatic variation and the geographical distribution of sex-determining mechanisms in the housefly. *Evolutionary Ecology Research*, 10(6), 797–809.
- Feller, A. F., Ogi, V., Seehausen, O., & Meier, J. I. (2021). Identification of a novel sex determining chromosome in cichlid fishes that acts as XY or ZW in different lineages. *Hydrobiologia*, 848(16), 3727–3745. <https://doi.org/10.1007/s10750-021-04560-7>
- Felsenstein, J. (1974). The evolutionary advantage of recombination. *Genetics*, 78(2), 737–756. <https://doi.org/10.1093/genetics/78.2.737>
- Fitzpatrick, J. L., Kahrl, A. F., & Snook, R. R. (2022). SpermTree, a species-level database of sperm morphology spanning the animal tree of life. *Scientific Data*, 9(1), 30. <https://doi.org/10.1038/s41597-022-01131-w>
- Foy, P. D., Loetzerich, S. R., Boxler, D., ... Meisel, R. P. (2024). Frequencies of house fly proto-Y chromosomes across populations are predicted by temperature heterogeneity within populations. *The Journal of Heredity*, esae056. <https://doi.org/10.1093/jhered/esae056>
- Fraïsse, C., Picard, M. A. L., & Vicoso, B. (2017). The deep conservation of the Lepidoptera Z chromosome suggests a non-canonical origin of the W. *Nature Communications*, 8(1), 1486. <https://doi.org/10.1038/s41467-017-01663-5>
- Fraïsse, C., & Sachdeva, H. (2021). The rates of introgression and barriers to genetic exchange between hybridizing species: Sex chromosomes vs autosomes. *Genetics*, 217(2), iyaa025. <https://doi.org/10.1093/genetics/iyaa025>
- Fredga, K. (1988). Aberrant chromosomal sex-determining mechanisms in mammals, with special reference to species with XY females. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 322(1208), 83–95. <https://doi.org/10.1098/rstb.1988.0116>
- Freitas, S., Parker, D. J., Labédan, M., ... Schwander, T. (2023). Evidence for cryptic sex in parthenogenetic stick insects of the genus *Timema*. *Proceedings Biological Sciences*, 290(2007), 20230404. <https://doi.org/10.1098/rspb.2023.0404>
- Fridolfsson, A. K., Cheng, H., Copeland, N. G., ... Ellegren, H. (1998). Evolution of the avian sex chromosomes from an ancestral pair of autosomes. *Proceedings of the National Academy of Sciences of the United States of America*, 95(14), 8147–8152. <https://doi.org/10.1073/pnas.95.14.8147>
- Froese, R., & Pauly, D. (2024). FishBase. [www.fishbase.org](http://www.fishbase.org)
- Fuad, M. M. H., Vetešník, L., & Šimková, A. (2021). Is gynogenetic reproduction in gibel carp (*Carassius gibelio*) a major trait responsible for invasiveness? *Journal of Vertebrate Biology*, 70(4), 21049.1–2104913.
- Gamble, T., Coryell, J., Ezaz, T., ... Zarkower, D. (2015). Restriction DNA Sequencing (RAD-seq) reveals an extraordinary number of transitions among gecko sex-determining systems. *Molecular Biology and Evolution*, 32(5), 1296–1309. <https://doi.org/10.1093/molbev/msv023>
- Garcia, S., Garnatje, T., & Kovařík, A. (2012). Plant rDNA database: Ribosomal DNA loci information goes online. *Chromosoma*, 121(4), 389–394. <https://doi.org/10.1007/s00412-012-0368-7>
- Garcia, S., Janousek, B., Pascual-Díaz, J. P., & Renner, S. S. (2023). Sex-chrom v. 2.0: A database of green plant species with sex chromosomes. *Chromosoma*, 132(1), 55–58. <https://doi.org/10.1007/s00412-023-00786-7>
- Ge, C., Ye, J., Weber, C., ... Capel, B. (2018). The histone demethylase KDM6B regulates temperature-dependent sex determination in a turtle species. *Science*, 360(6389), 645–648. <https://doi.org/10.1126/science.aap8328>
- Ghiselin, M. (1969). The evolution of hermaphroditism among animals. *The Quarterly Review of Biology*, 44(2), 189–208.
- Gil-Fernández, A., Saunders, P. A., Martín-Ruiz, M., ... Page, J. (2020). Meiosis reveals the early steps in the evolution of a neo-XY sex chromosome pair in the African pygmy mouse *Mus minutoides*. *PLoS Genetics*, 16(11), e1008959. <https://doi.org/10.1371/journal.pgen.1008959>
- Godin, V. N. (2022). Trioecy in flowering plants. *Doklady Biological Sciences*, 507(1), 301–311. <https://doi.org/10.1134/S0012496622060023>
- Goldberg, E. E., Kohn, J. R., Lande, R., ... Igić, B. (2010). Species selection maintains self-incompatibility. *Science (New York, N.Y.)*, 330(6003), 493–495. <https://doi.org/10.1126/science.1194513>
- Goldberg, E. E., Otto, S. P., Vamasi, J. C., ... Ashman, T. -L. (2017). Macroevolutionary synthesis of flowering plant sexual systems. *Evolution; International Journal of Organic Evolution*, 71(4), 898–912. <https://doi.org/10.1111/evo.13181>
- Gómez, J. M., González-Megías, A., & Verdú, M. (2023). The evolution of same-sex sexual behaviour in mammals. *Nature Communications*, 14(1), 5719. <https://doi.org/10.1038/s41467-023-41290-x>
- Goodwillie, C., Kalisz, S., & Eckert, C. (2005a). The evolutionary enigma of mixed mating systems in plants: Occurrence, theoretical explanations, and empirical evidence. *Annual Review of Ecology, Evolution, and Systematics*, 36, 47–79.
- Goodwillie, C., Kalisz, S., & Eckert, C. G. (2005b). The evolutionary enigma of mixed mating systems in plants: Occurrence, theoretical explanations, and empirical evidence. *Annual Review of Ecology, Evolution, and Systematics*, 36(1), 47–79. <https://doi.org/10.1146/annurev.ecolsys.36.091704.175539>
- Goodwillie, C., Sargent, R. D., Eckert, C. G., ... Winn, A. A. (2010). Correlated evolution of mating system and floral display traits in flowering plants and its implications for the distribution of mating system variation. *The New Phytologist*, 185(1), 311–321. <https://doi.org/10.1111/j.1469-8137.2009.03043.x>
- Graves, J. A. (1995). The origin and function of the mammalian Y chromosome and Y-borne genes – An evolving understanding. *BioEssays*, 17(4), 311–320. <https://doi.org/10.1002/bies.950170407>
- Green, R. F., & Noakes, D. L. G. (1995). Is a little bit of sex as good as a lot? *Journal of Theoretical Biology*, 174(1), 87–96.
- Gregory, T. R. (2024). *Animal genome size database*. [www.genomesize.com](http://www.genomesize.com)
- Grigoriev, I. V., Hayes, R. D., Calhoun, S., ... Kuo, A. (2021). PhycoCosm, a comparative algal genomics resource. *Nucleic Acids Research*, 49(D1), D1004–D1011. <https://doi.org/10.1093/nar/gkaa898>
- Grossenbacher, D., Briscoe Runquist, R., Goldberg, E. E., & Brandvain, Y. (2015). Geographic range size is predicted by plant mating system. *Ecology Letters*, 18(7), 706–713. <https://doi.org/10.1111/ele.12449>
- Gutiérrez, M. L., Rodríguez-González, R., Fuentes, I., ... Garcia, S. (2023). First update to B-chrom: A database on B-chromosomes. *Methods in Molecular Biology (Clifton, N.J.)*, 2703, 227–236.
- Haldane, J. B. S. (1922). Sex ratio and unisexual sterility in hybrid animals. *Journal of Genetics*, 12(2), 101–109. <https://doi.org/10.1007/bf02983075>

- Hamilton, W. D. (1967). Extraordinary sex ratios. A sex-ratio theory for sex linkage and inbreeding has new implications in cytogenetics and entomology. *Science (New York, N.Y.)*, 156(3774), 477–488. <https://doi.org/10.1126/science.156.3774.477>
- Hargreaves, A. L., & Eckert, C. G. (2014). Evolution of dispersal and mating systems along geographic gradients: Implications for shifting ranges. *Functional Ecology*, 28(1), 5–21. <https://doi.org/10.1111/1365-2435.12170>
- Hartfield, M. (2016). Evolutionary genetic consequences of facultative sex and outcrossing. *Journal of Evolutionary Biology*, 29(1), 5–22. <https://doi.org/10.1111/jeb.12770>
- Hartfield, M., Bataillon, T., & Glémin, S. (2017). The evolutionary interplay between adaptation and self-fertilization. *Trends in Genetics: TIG*, 33(6), 420–431. <https://doi.org/10.1016/j.tig.2017.04.002>
- Hartfield, M., & Keightley, P. D. (2012). Current hypotheses for the evolution of sex and recombination. *Integrative Zoology*, 7(2), 192–209. <https://doi.org/10.1111/j.1749-4877.2012.00284.x>
- Hedrick, P. W. (1987). Population genetics of intragametophytic selfing. *Evolution; International Journal of Organic Evolution*, 41(1), 137–144. <https://doi.org/10.1111/j.1558-5646.1987.tb05776.x>
- Heesch, S., Serrano-Serrano, M., Barrera-Redondo, J., ... Coelho, S. M. (2021). Evolution of life cycles and reproductive traits: Insights from the brown algae. *Journal of Evolutionary Biology*, 34(7), 992–1009. <https://doi.org/10.1111/jeb.13880>
- Hill, W. G., & Robertson, A. (1966). The effect of linkage on limits to artificial selection. *Genetical Research*, 8(3), 269–294.
- Hodson, C. N., & Ross, L. (2021). Evolutionary perspectives on germline-restricted chromosomes in flies (Diptera). *Genome Biology and Evolution*, 13(6), evab072. <https://doi.org/10.1093/gbe/evab072>
- Hoencamp, C., Dudchenko, O., Elbarsh, A. M. O., ... Rowland, B. D. (2021). 3D genomics across the tree of life reveals condensin II as a determinant of architecture type. *Science*, 372(6545), 984–989. <https://doi.org/10.1126/science.abe2218>
- Hofstatter, P. G., & Lahr, D. J. G. (2019). All eukaryotes are sexual, unless proven otherwise: Many so-called asexuals present meiotic machinery and might be able to have sex. *BioEssays*, 41(6), e1800246. <https://doi.org/10.1002/bies.201800246>
- Holleley, C. E., O'Meally, D., Sarre, S. D., ... Georges, A. (2015). Sex reversal triggers the rapid transition from genetic to temperature-dependent sex. *Nature*, 523(7558), 79–82. <https://doi.org/10.1038/nature14574>
- Holleley, C. E., Sarre, S. D., O'Meally, D., & Georges, A. (2016). Sex reversal in reptiles: Reproductive oddity or powerful driver of evolutionary change? *Sexual Development: Genetics, Molecular Biology, Evolution, Endocrinology, Embryology, and Pathology of Sex Determination and Differentiation*, 10(5-6), 279–287. <https://doi.org/10.1159/000450972>
- Honeycutt, J., Deck, C., Miller, S. C., & Godwin, J. (2019). Warmer waters masculinize wild populations of a fish with temperature-dependent sex determination. *Scientific Reports*, 9, 6527. <https://doi.org/10.1038/s41598-019-42944-x>
- Hood, M. E., & Antonovics, J. (2004). Mating within the meiotic tetrad and the maintenance of genomic heterozygosity. *Genetics*, 166(4), 1751–1759. <https://doi.org/10.1534/genetics.166.4.1751>
- Hughes, J. F., & Page, D. C. (2015). The biology and evolution of mammalian Y chromosomes. *Annual Review of Genetics*, 49(1), 507–527. <https://doi.org/10.1146/annurev-genet-112414-055311>
- Hughes, J. J., Lagunas-Robles, G., Campbell, P., & Fishman, L. (2024). The role of conflict in the formation and maintenance of variant sex chromosome systems in mammals. *Journal of Heredity*, 115(6), 601–624. <https://doi.org/10.1093/jhered/esae031>
- Imarazene, B., Du, K., Beille, S., ... Guiguen, Y. (2021). A supernumerary “B-sex” chromosome drives male sex determination in the Pachón cavefish, *Astyanax mexicanus*. *Current Biology*, 31(21), 4800–4809.e9. <https://doi.org/10.1016/j.cub.2021.08.030>
- Immler, S., & Otto, S. P. (2015). The evolution of sex chromosomes in organisms with separate haploid sexes. *Evolution; International Journal of Organic Evolution*, 69(3), 694–708. <https://doi.org/10.1111/evo.12602>
- Ioannidis, J., Taylor, G., Zhao, D., ... Clinton, M. (2021). Primary sex determination in birds depends on DMRT1 dosage, but gonadal sex does not determine adult secondary sex characteristics. *Proceedings of the National Academy of Sciences of the United States of America*, 118(10), e2020909118. <https://doi.org/10.1073/pnas.2020909118>
- Iwasaki, M., Kajiwar, T., Yasui, Y., ... Kohchi, T. (2021). Identification of the sex-determining factor in the liverwort *Marchantia polymorpha* reveals unique evolution of sex chromosomes in a haploid system. *Current Biology: CB*, 31(24), 5522–5532.e7. <https://doi.org/10.1016/j.cub.2021.10.023>
- Janicke, T., Häderer, I. K., Lajeunesse, M. J., & Anthes, N. (2016). Darwinian sex roles confirmed across the animal kingdom. *Science Advances*, 2(2), e1500983. <https://doi.org/10.1126/sciadv.1500983>
- Jankásek, M., Kotyková Varadinová, Z., & Štáhlavský, F. (2021). Blatodea karyotype database. *European Journal of Entomology*, 118, 192–199. <https://doi.org/10.14411/eje.2021.020>
- Janko, K., Bartoš, O., Kočí, J., ... Štefková-Kašparová, E. (2021). Genome fractionation and loss of heterozygosity in hybrids and polyploids: Mechanisms, consequences for selection, and link to gene function. *Molecular Biology and Evolution*, 38(12), 5255–5274. <https://doi.org/10.1093/molbev/msab249>
- Jarne, P., & Auld, J. R. (2006). Animals mix it up too: The distribution of self-fertilization among hermaphroditic animals. *Evolution; International Journal of Organic Evolution*, 60(9), 1816–1824. <https://doi.org/10.1554/06-246.1>
- Jarne, P., & Charlesworth, D. (1993). The evolution of the selfing rate in functionally hermaphrodite plants and animals. *Annual Review of Ecology, Evolution, and Systematics*, 24, 441–466.
- Jaron, K. S., Bast, J., Nowell, R. W., ... Schwander, T. (2021). Genomic features of parthenogenetic animals. *The Journal of Heredity*, 112(1), 19–33. <https://doi.org/10.1093/jhered/esaa031>
- Jaron, K. S., Parker, D. J., Anselmetti, Y., ... Schwander, T. (2022). Convergent consequences of parthenogenesis on stick insect genomes. *Science Advances*, 8(8), eabg3842. <https://doi.org/10.1126/sciadv.abg3842>
- Jay, P., Jeffries, D., Hartmann, F. E., ... Giraud, T. (2024). Why do sex chromosomes progressively lose recombination? *Trends in Genetics: TIG*, 40(7), 564–579. <https://doi.org/10.1016/j.tig.2024.03.005>
- Jeffries, D. L., Gerchen, J. F., Scharmann, M., & Pannell, J. R. (2021). A neutral model for the loss of recombination on sex chromosomes. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 376(1832), 20200096. <https://doi.org/10.1098/rstb.2020.0096>
- Jeffries, D. L., Lavanchy, G., Sermier, R., ... Perrin, N. (2018). A rapid rate of sex-chromosome turnover and non-random transitions in true frogs. *Nature Communications*, 9(1), 4088. <https://doi.org/10.1038/s41467-018-06517-2>
- Jensen, M. P., Allen, C. D., Eguchi, T., ... Dutton, P. H. (2018). Environmental warming and feminization of one of the largest sea turtle populations in the world. *Current Biology: CB*, 28(1), 154–159.e4. <https://doi.org/10.1016/j.cub.2017.11.057>
- Jerkovic, I., & Cavalli, G. (2021). Understanding 3D genome organization by multidisciplinary methods. *Nature Reviews. Molecular cell biology*, 22(8), 511–528. <https://doi.org/10.1038/s41580-021-00362-w>
- Jesson, L. K., & Garnock-Jones, P. J. (2011). Can classifications of functional gender be extended to all land plants? *Perspectives in Plant Ecology, Evolution and Systematics*, 14(2), 153–160. <https://doi.org/10.1016/j.ppees.2011.10.003>
- Johnson Pokorná, M., & Kratochvíl, L. (2016). What was the ancestral sex-determining mechanism in amniote vertebrates? *Biological Reviews of the Cambridge Philosophical Society*, 91(1), 1–12. <https://doi.org/10.1111/brv.12156>
- Jonika, M. M., Alfieri, J. M., Sylvester, T., ... Blackmon, H. (2022). Why not Y naught. *Heredity*, 129(2), 75–78. <https://doi.org/10.1038/s41437-022-00543-z>
- Joshi, C. H., & Wiens, J. J. (2023). Does haplodiploidy help drive the evolution of insect eusociality? *Frontiers in Ecology and Evolution*, 11, 11. <https://doi.org/10.3389/fevo.2023.1118748>

- Käfer, J. (2022). How sex chromosomes get trapped into nonrecombination. *PLoS Biology*, 20(7), e3001718. <https://doi.org/10.1371/journal.pbio.3001718>
- Käfer, J., Marais, G. A. B., & Pannell, J. R. (2017). On the rarity of dioecy in flowering plants. *Molecular Ecology*, 26(5), 1225–1241. <https://doi.org/10.1111/mec.14020>
- Kato, Y., Nitta, J. H., Perez, C. A. G., ... Watanabe, H. (2024). Identification of gene isoforms and their switching events between male and female embryos of the parthenogenetic crustacean *Daphnia magna*. *Scientific Reports*, 14(1), 9407. <https://doi.org/10.1038/s41598-024-59774-1>
- Kent, T. V., Uzunović, J., & Wright, S. I. (2017). Coevolution between transposable elements and recombination. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 372(1736), 20160458. <https://doi.org/10.1098/rstb.2016.0458>
- Khalaf, A., Francis, O., & Blaxter, M. L. (2024). Genome evolution in intracellular parasites: Microsporidia and Apicomplexa. *The Journal of Eukaryotic Microbiology*, 71(5), e13033. <https://doi.org/10.1111/jeu.13033>
- Kitano, J., Ansai, S., Takehana, Y., & Yamamoto, Y. (2024). Diversity and convergence of sex-determination mechanisms in teleost fish. *Annual Review of Animal Biosciences*, 12, 233–259. <https://doi.org/10.1146/annurev-animal-021122-113935>
- Kocher, T. D., Meisel, R. P., Gamble, T., ... Gammerdinger, W. J. (2024). Yes, polygenic sex determination is a thing! *Trends in Genetics: TIG*, 40(12), 1001–1017. <https://doi.org/10.1016/j.tig.2024.10.003>
- Kočí, J., Röslein, J., Pačes, J., ... Janko, K. (2020). No evidence for accumulation of deleterious mutations and fitness degradation in clonal fish hybrids: Abandoning sex without regrets. *Molecular Ecology*, 29(16), 3038–3055. <https://doi.org/10.1111/mec.15539>
- Komen, H., & Thorgaard, G. H. (2007). Androgenesis, gynogenesis and the production of clones in fishes: A review. *Aquaculture*, 269(1–4), 150–173. <https://doi.org/10.1016/j.aquaculture.2007.05.009>
- Kořínková, T., & Král, J. (2013). Karyotypes, sex chromosomes, and meiotic division in spiders. In Nentwig, W. (Ed.), *Spider ecophysiology* (pp. 159–171). Springer Berlin Heidelberg.
- Kratochvíl, L., Gamble, T., & Rovatsos, M. (2021a). Sex chromosome evolution among amniotes: Is the origin of sex chromosomes non-random? *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 376(1833), 20200108. <https://doi.org/10.1098/rstb.2020.0108>
- Kratochvíl, L., Stöck, M., Rovatsos, M., ... Pokorná, M. J. (2021b). Expanding the classical paradigm: What we have learnt from vertebrates about sex chromosome evolution. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 376(1833), 20200097. <https://doi.org/10.1098/rstb.2020.0097>
- Krueger, C. J., & Janzen, F. J. (2022). ROSIE, a database of reptilian offspring sex ratios and sex-determining mechanisms, beginning with Testudines. *Scientific Data*, 9(1), 22. <https://doi.org/10.1038/s41597-021-01108-1>
- Krueger-Hadfield, S. A. (2024). Let's talk about sex: Why reproductive systems matter for understanding algae. *Journal of Phycology*, 60(3), 581–597. <https://doi.org/10.1111/jpy.13462>
- Krueger-Hadfield, S. A., Shainker-Connolly, S. J., Crowell, R. M., & Vis, M. L. (2024). The eco-evolutionary importance of reproductive system variation in the macroalgae: Freshwater reds as a case study. *Journal of Phycology*, 60(1), 15–25. <https://doi.org/10.1111/jpy.13407>
- Kuhl, H., Guiguen, Y., Höhne, C., ... Stöck, M. (2021). A 180 Myr-old female-specific genome region in sturgeon reveals the oldest known vertebrate sex determining system with undifferentiated sex chromosomes. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 376(1832), 20200089. <https://doi.org/10.1098/rstb.2020.0089>
- Lampert, K. P., & Scharl, M. (2008). The origin and evolution of a unisexual hybrid: *Poecilia formosa*. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 363(1505), 2901–2909. <https://doi.org/10.1098/rstb.2008.0040>
- Lee, S. C., Corradi, N., Byrnes, E. J. 3rd, ... Heitman, J. (2008). Microsporidia evolved from ancestral sexual fungi. *Current Biology: CB*, 18(21), 1675–1679. <https://doi.org/10.1016/j.cub.2008.09.030>
- Lehtonen, J., Horinouchi, Y., Togashi, T., & Parker, G. A. (2021). Evolution of anisogamy in organisms with parthenogenetic gametes. *The American Naturalist*, 198(3), 360–378. <https://doi.org/10.1086/715185>
- Lehtonen, J., & Kokko, H. (2011). Two roads to two sexes: Unifying gamete competition and gamete limitation in a single model of anisogamy evolution. *Behavioral Ecology and Sociobiology*, 65(3), 445–459. <https://doi.org/10.1007/s00265-010-1116-8>
- Lehtonen, J., Kokko, H., & Parker, G. A. (2016). What do isogamous organisms teach us about sex and the two sexes? *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 371(1706), 20150532. <https://doi.org/10.1098/rstb.2015.0532>
- Lehtonen, J., & Parker, G. A. (2014). Gamete competition, gamete limitation, and the evolution of the two sexes. *Molecular Human Reproduction*, 20(12), 1161–1168. <https://doi.org/10.1093/molehr/gau068>
- Leitch, I. J., Johnston, E., Pellicer, J., Hidalgo, O., & Bennett. (2019). *Plant DNA C-values Database, Release 7.1*. <https://cvalues.science.kew.org/>
- Lenormand, T., & Roze, D. (2024). A single theory for the evolution of sex chromosomes and the two rules of speciation. *bioRxiv* 585601, <https://doi.org/10.1101/2024.03.18.585601>, March 20, 2024, preprint: not peer reviewed.
- Leonard, J. L. (Ed.). (2019). *Transitions between sexual systems: Understanding the mechanisms of, and pathways between, dioecy, hermaphroditism and other sexual systems* (1st ed.). Springer International Publishing.
- Leung, K., & van der Meulen, H. (2022). Revisiting the hymenopteran diploid male vortex: A review of avoidance mechanisms and incidence. *Entomologia Experimentalis et Applicata*, 170(12), 1010–1031. <https://doi.org/10.1111/eea.13234>
- Li, J. (2022). A review of sexual determination and differentiation in crustacean. *Journal of Biosciences and Medicines*, 10(07), 19–37. <https://doi.org/10.4236/jbm.2022.107002>
- Li, X., -Y., Liu, X. -L., Zhu, Y. -J., ... Gui, J. -F. (2018). Origin and transition of sex determination mechanisms in a gynogenetic hexaploid fish. *Heredity*, 121(1), 64–74. <https://doi.org/10.1038/s41437-017-0049-7>
- Li, Y., Chen, Z., Liu, H., ... Luo, D. (2021). ASER: Animal sex reversal database. *Genomics, Proteomics & Bioinformatics*, 19(6), 873–881. <https://doi.org/10.1016/j.gpb.2021.10.001>
- Liehr, T. (2021). Molecular cytogenetics in the era of chromosomics and cytogenomic approaches. *Frontiers in Genetics*, 12, 720507. <https://doi.org/10.3389/fgene.2021.720507>
- Lima, T. G. (2014). Higher levels of sex chromosome heteromorphism are associated with markedly stronger reproductive isolation. *Nature Communications*, 5(1), 1–8.
- Linck, E. B., & Cadena, C. D. (2024). A latitudinal gradient of reference genomes. *bioRxiv* 602657. <https://doi.org/10.1101/2024.07.09.602657>, July 13, 2024, preprint: not peer reviewed.
- Lipinska, A. P., Cossard, G., Epperlein, P., ... Coelho, S. M. (2024). Structural and evolutionary features of red algal UV sex chromosomes. *bioRxiv* 626989. <https://doi.org/10.1101/2024.12.05.626989>, December 09, 2024, preprint: not peer reviewed.
- Lisachov, A. P., Tishakova, K. V., Romanenko, S. A., ... Trifonov, V. A. (2021). Whole-chromosome fusions in the karyotype evolution of *Sceloporus* (Iguania, Reptilia) are more frequent in sex chromosomes than autosomes. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 376(1833), 20200099. <https://doi.org/10.1098/rstb.2020.0099>
- Liu, P., Vigneau, J., Craig, R. J., ... Coelho, S. M. (2024). 3D chromatin maps of a brown alga reveal U/V sex chromosome spatial organization. *Nature Communications*, 15(1), 9590. <https://doi.org/10.1038/s41467-024-53453-5>

- Lockley, E. C., & Eizaguirre, C. (2021). Effects of global warming on species with temperature-dependent sex determination: Bridging the gap between empirical research and management. *Evolutionary Applications*, 14(10), 2361–2377. <https://doi.org/10.1111/eva.13226>
- Ma, W. -J., & Veltsos, P. (2021). The diversity and evolution of sex chromosomes in frogs. *Genes*, 12(4), 483. <https://doi.org/10.3390/genes12040483>
- Marie-Orleach, L., Brochmann, C., & Glémin, S. (2022). Mating system and speciation I: Accumulation of genetic incompatibilities in allopatry. *PLoS Genetics*, 18(12), e1010353. <https://doi.org/10.1371/journal.pgen.1010353>
- Mark Welch, D. B., Mark Welch, J. L., & Meselson, M. (2008). Evidence for degenerate tetraploidy in bdelloid rotifers. *Proceedings of the National Academy of Sciences of the United States of America*, 105(13), 5145–5149. <https://doi.org/10.1073/pnas.0800972105>
- Masly, J. P., & Presgraves, D. C. (2007). High-resolution genome-wide dissection of the two rules of speciation in *Drosophila*. *PLoS Biology*, 5(9), e243. <https://doi.org/10.1371/journal.pbio.0050243>
- Matsumoto, T., & Kitano, J. (2016). The intricate relationship between sexually antagonistic selection and the evolution of sex chromosome fusions. *Journal of Theoretical Biology*, 404, 97–108. <https://doi.org/10.1016/j.jtbi.2016.05.036>
- McCauley, D. E., & Bailey, M. F. (2009). Recent advances in the study of gynodioecy: The interface of theory and empiricism. *Annals of Botany*, 104(4), 611–620. <https://doi.org/10.1093/aob/mcp141>
- Meyer, B. J. (2022). Mechanisms of sex determination and X-chromosome dosage compensation. *Genetics*, 220(2), iyab197. <https://doi.org/10.1093/genetics/iyab197>
- Ming, R., Bendahmane, A., & Renner, S. S. (2011). Sex chromosomes in land plants. *Annual Review of Plant Biology*, 62, 485–514. <https://doi.org/10.1146/annurev-arplant-042110-103914>
- Mirzaghaderi, G., & Hörandl, E. (2016). The evolution of meiotic sex and its alternatives. *Proceedings. Biological sciences/The Royal Society*, 283(1838), 20161221. <https://doi.org/10.1098/rspb.2016.1221>
- Mizoguchi, B. A., & Valenzuela, N. (2016). Ecotoxicological perspectives of sex determination. *Sexual Development*, 10(1), 45–57. <https://doi.org/10.1159/000444770>
- Mogensen, G. S. (1985). *Illustrated moss flora of Arctic North America and Greenland* (Vol. 17). Museum Tusculanum Press.
- Morelli, M. W., Blackmon, H., & Hjelman, C. E. (2022). Diptera and *Drosophila* karyotype databases: A useful dataset to guide evolutionary and genomic studies. *Frontiers in Ecology and Evolution*, 10, 10. <https://doi.org/10.3389/fevo.2022.832378>
- Morgan, T. H. (1914). No crossing over in the male of *Drosophila* of genes in the second and third pairs of chromosomes. *The Biological Bulletin*, 26(4), 195–204. <https://doi.org/10.2307/1536193>
- Morgan-Richards, M., Treweek, S. A., & Stringer, I. A. N. (2010). Geographic parthenogenesis and the common tea-tree stick insect of New Zealand. *Molecular Ecology*, 19(6), 1227–1238. <https://doi.org/10.1111/j.1365-294x.2010.04542.x>
- Mork, L., Czerwinski, M., & Capel, B. (2014). Predetermination of sexual fate in a turtle with temperature-dependent sex determination. *Developmental Biology*, 386(1), 264–271. <https://doi.org/10.1016/j.ydbio.2013.11.026>
- Muller, H. J. (1964). The relation of recombination to mutational advance. *Mutation Research*, 106, 2–9. [https://doi.org/10.1016/0027-5107\(64\)90047-8](https://doi.org/10.1016/0027-5107(64)90047-8)
- Mulugeta, E., Wassenaar, E., Sleddens-Linkels, E., ... Baarends, W. M. (2016). Genomes of Ellobius species provide insight into the evolutionary dynamics of mammalian sex chromosomes. *Genome Research*, 26(9), 1202–1210.
- Myosho, T., Takehana, Y., Hamaguchi, S., & Sakaizumi, M. (2015). Turnover of sex chromosomes in celebensis group medaka fishes. *G3 (Bethesda, Md.)*, 5(12), 2685–2691. <https://doi.org/10.1534/g3.115.021543>
- Nagpure, N. S., Pathak, A. K., Pati, R., ... Murali, S. (2016). Fish Karyome version 2.1: A chromosome database of fishes and other aquatic organisms. *Database*, 2016, baw012. <https://doi.org/10.1093/database/baw012>
- Neaves, W. B., & Baumann, P. (2011). Unisexual reproduction among vertebrates. *Trends in Genetics: TIG*, 27(3), 81–88. <https://doi.org/10.1016/j.tig.2010.12.002>
- Neiman, M., Meirmans, P. G., Schwander, T., & Meirmans, S. (2018). Sex in the wild: How and why field-based studies contribute to solving the problem of sex. *Evolution; International Journal of Organic Evolution*, 72(6), 1194–1203. <https://doi.org/10.1111/evo.13485>
- Neiman, M., Sharbel, T. F., & Schwander, T. (2014). Genetic causes of transitions from sexual reproduction to asexuality in plants and animals. *Journal of Evolutionary Biology*, 27(7), 1346–1359. <https://doi.org/10.1111/jeb.12357>
- Nemesházi, E., & Bókony, V. (2022). Asymmetrical sex reversal: Does the type of heterogamety predict propensity for sex reversal? *Bio-Essays*, 44(7), e2200039. <https://doi.org/10.1002/bies.202200039>
- Nemesházi, E., & Bókony, V. (2023). HerpSexDet: The herpetological database of sex determination and sex reversal. *Scientific Data*, 10(1), 377. <https://doi.org/10.1038/s41597-023-02268-y>
- Nemesházi, E., Gál, Z., Ujhegyi, N., ... Bókony, V. (2020). Novel genetic sex markers reveal high frequency of sex reversal in wild populations of the agile frog (*Rana dalmatina*) associated with anthropogenic land use. *Molecular Ecology*, 29(19), 3607–3621. <https://doi.org/10.1111/mec.15596>
- Nemesházi, E., Kövér, S., & Bókony, V. (2021). Evolutionary and demographic consequences of temperature-induced masculinization under climate warming: the effects of mate choice. *BMC Ecology and Evolution*, 21(1), 16. <https://doi.org/10.1186/s12862-021-01747-3>
- Nguyen, A. H., & Bachtrog, D. (2021). Toxic Y chromosome: Increased repeat expression and age-associated heterochromatin loss in male *Drosophila* with a young Y chromosome. *PLoS Genetics*, 17(4), e1009438. <https://doi.org/10.1371/journal.pgen.1009438>
- Nieuwenhuis, B. P. S., & James, T. Y. (2016). The frequency of sex in fungi. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 371(1706), 20150540. <https://doi.org/10.1098/rstb.2015.0540>
- Nigon, V. (1949). Les modalités de la reproduction et le déterminisme du sexe chez quelques nématodes libres. *Annales des Sciences Naturelles - Zoologie et Biologie Animale*, 2, 1–132.
- Normark, B. B. (2003). The evolution of alternative genetic systems in insects. *Annual Review of Entomology*, 48(1), 397–423. <https://doi.org/10.1146/annurev.ento.48.091801.112703>
- Normark, B. B. (2004). Haplodiploidy as an outcome of coevolution between male-killing cytoplasmic elements and their hosts. *Evolution; International Journal of Organic Evolution*, 58(4), 790–798. <https://doi.org/10.1111/j.0014-3820.2004.tb00412.x>
- Núñez, O. (1962). Cytology of collembola. *Nature*, 194(4832), 946–947. <https://doi.org/10.1038/194946a0>
- Okada, S., Sone, T., Fujisawa, M., ... Ohya, K. (2001). The Y chromosome in the liverwort *Marchantia polymorpha* has accumulated unique repeat sequences harboring a male-specific gene. *Proceedings of the National Academy of Sciences of the United States of America*, 98(16), 9454–9459. <https://doi.org/10.1073/pnas.171304798>
- Otto, S. P. (2009). The evolutionary enigma of sex. *The American Naturalist*, 174(Suppl 1), S1–S14. <https://doi.org/10.1086/599084>
- Otto, S. P. (2021). Selective interference and the evolution of sex. *The Journal of Heredity*, 112(1), 9–18. <https://doi.org/10.1093/jhered/esaa026>
- Oyarzún, P. A., Nuñez, J. J., Toro, J. E., & Gardner, J. P. A. (2020). Trioecy in the marine mussel *Semimytilus algosus* (Mollusca, Bivalvia): Stable sex ratios across 22 degrees of a latitudinal gradient. *Frontiers in Marine Science*, 7, 7. <https://doi.org/10.3389/fmars.2020.00348>
- Öztoprak, H., Gao, S., Guiguelmoni, N., ... Bast, J. (2023). Haplotype independence contributes to evolvability in the long-term absence of sex in a mite. *bioRxiv* 556471.

- <https://doi.org/10.1101/2023.09.07.556471>, October 20, 2023, preprint: not peer reviewed.
- Pan, Q., Feron, R., Yano, A., ... Guiguen, Y. (2019). Identification of the master sex determining gene in Northern pike (*Esox lucius*) reveals restricted sex chromosome differentiation. *PLoS Genetics*, 15(8), e1008013. <https://doi.org/10.1371/journal.pgen.1008013>
- Pannell, J. R. (2015). Evolution of the mating system in colonizing plants. *Molecular Ecology*, 24(9), 2018–2037. <https://doi.org/10.1111/mec.13087>
- Pannell, J. R., Auld, J. R., Brandvain, Y., ... Winn, A. A. (2015). The scope of Baker's law. *The New Phytologist*, 208(3), 656–667. <https://doi.org/10.1111/nph.13539>
- Pannell, J. R., & Jordan, C. Y. (2022). Evolutionary transitions between hermaphroditism and dioecy in animals and plants. *Annual Review of Ecology, Evolution, and Systematics*, 53(1), 183–201. <https://doi.org/10.1146/annurev-ecolsys-102320-085812>
- Parker, G. A. (1978). Selection on non-random fusion of gametes during the evolution of anisogamy. *Journal of Theoretical Biology*, 73(1), 1–28. [https://doi.org/10.1016/0022-5193\(78\)90177-7](https://doi.org/10.1016/0022-5193(78)90177-7)
- Pellicer, J., & Leitch, I. J. (2020). The plant DNA C-values database (release 7.1): An updated online repository of plant genome size data for comparative studies. *The New Phytologist*, 226(2), 301–305. <https://doi.org/10.1111/nph.16261>
- Pennell, M. W., Kirkpatrick, M., Otto, S. P., ... Kitano, J. (2015). Y fuse? Sex chromosome fusions in fishes and reptiles. *PLoS Genetics*, 11(5), e1005237. <https://doi.org/10.1371/journal.pgen.1005237>
- Pennell, M. W., Mank, J. E., & Peichel, C. L. (2018). Transitions in sex determination and sex chromosomes across vertebrate species. *Molecular Ecology*, 27(19), 3950–3963. <https://doi.org/10.1111/mec.14540>
- Peona, V., Palacios-Gimenez, O. M., Blommaert, J., ... Suh, A. (2021). The avian W chromosome is a refuge for endogenous retroviruses with likely effects on female-biased mutational load and genetic incompatibilities. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 376(1833), 20200186. <https://doi.org/10.1098/rstb.2020.0186>
- Perkins, R. D., Gamboa, J. R., Jonika, M. M., ... Blackmon, H. (2019). A database of amphibian karyotypes. *Chromosome Research*, 27(4), 313–319. <https://doi.org/10.1007/s10577-019-09613-1>
- Perrin, N. (2016). Random sex determination: When developmental noise tips the sex balance. *BioEssays*, 38(12), 1218–1226. <https://doi.org/10.1002/bies.201600093>
- The Phaeoexplorer project. (2024). *Phaeoexplorer brown algal genome database*. <https://phaeoexplorer.sb-roscoff.fr/about/>
- Phillips, B. C., Rodrigues, N., Jansen van Rensburg, A., & Perrin, N. (2020). Phylogeography, more than elevation, accounts for sex chromosome differentiation in Swiss populations of the common frog (*Rana temporaria*). *Evolution; International Journal of Organic Evolution*, 74(3), 644–654. <https://doi.org/10.1111/evo.13860>
- Pickup, M., Brandvain, Y., Fraïsse, C., ... Field, D. L. (2019). Mating system variation in hybrid zones: Facilitation, barriers and asymmetries to gene flow. *The New Phytologist*, 224(3), 1035–1047. <https://doi.org/10.1111/nph.16180>
- Pickup, M., Brandvain, Y., Fraïsse, C., ... Field, D. L. (2020). *Data from: Mating system variation in hybrid zones: Facilitation, barriers and asymmetries to gene flow*. Dryad. <https://doi.org/10.5061/DRYAD.DM7CS86>
- Piferrer, F., Blázquez, M., Navarro, L., & González, A. (2005). Genetic, endocrine, and environmental components of sex determination and differentiation in the European sea bass (*Dicentrarchus labrax* L.). *General and Comparative Endocrinology*, 142(1–2), 102–110. <https://doi.org/10.1016/j.ygcen.2005.02.011>
- Pinto, B. J., Gamble, T., Smith, C. H., & Wilson, M. A. (2023). A lizard is never late: Squamate genomics as a recent catalyst for understanding sex chromosome and microchromosome evolution. *The Journal of Heredity*, 114(5), 445–458. <https://doi.org/10.1093/jhered/esad023>
- Pinto, B. J., Keating, S. E., Nielsen, S. V., ... Gamble, T. (2022). Chromosome-level genome assembly reveals dynamic sex chromosomes in neotropical leaf-litter geckos (Sphaerodactylidae: *Sphaerodactylus*). *The Journal of Heredity*, 113(3), 272–287. <https://doi.org/10.1093/jhered/esac016>
- Pla, S., Benvenuto, C., Capellini, I., & Piferrer, F. (2022). Switches, stability and reversals in the evolutionary history of sexual systems in fish. *Nature Communications*, 13(1), 3029. <https://doi.org/10.1038/s41467-022-30419-z>
- Pokorná, M., Altmanová, M., & Kratochvíl, L. (2014). Multiple sex chromosomes in the light of female meiotic drive in amniote vertebrates. *Chromosome Research*, 22(1), 35–44. <https://doi.org/10.1007/s10577-014-9403-2>
- Ponnikas, S., Sigeman, H., Abbott, J. K., & Hansson, B. (2018). Why do sex chromosomes stop recombining? *Trends in Genetics: TIG*, 34(7), 492–503. <https://doi.org/10.1016/j.tig.2018.04.001>
- Ramesh, M. A., Malik, S. -B., & Logsdon, J. M. Jr (2005). A phylogenomic inventory of meiotic genes; evidence for sex in *Giardia* and an early eukaryotic origin of meiosis. *Current Biology: CB*, 15(2), 185–191. <https://doi.org/10.1016/j.cub.2005.01.003>
- Rao, S. S. P., Huntley, M. H., Durand, N. C., ... Aiden, E. L. (2014). A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping. *Cell*, 159(7), 1665–1680.
- Razanajatovo, M., Maurel, N., Dawson, W., ... van Kleunen, M. (2016). Plants capable of selfing are more likely to become naturalized. *Nature Communications*, 7, 13313. <https://doi.org/10.1038/ncomms13313>
- Renner, S. S. (2014). The relative and absolute frequencies of angiosperm sexual systems: Dioecy, monoecy, gynodioecy, and an updated online database. *American Journal of Botany*, 101(10), 1588–1596. <https://doi.org/10.3732/ajb.1400196>
- Renner, S. S., & Müller, N. A. (2021). Plant sex chromosomes defy evolutionary models of expanding recombination suppression and genetic degeneration. *Nature Plants*, 7(4), 392–402. <https://doi.org/10.1038/s41477-021-00884-3>
- Rens, W., O'Brien, P. C. M., Grützner, F., ... Ferguson-Smith, M. A. (2007). The multiple sex chromosomes of platypus and echidna are not completely identical and several share homology with the avian Z. *Genome Biology*, 8(11), R243.
- Rice, A., Glick, L., Abadi, S., ... Mayrose, I. (2015). The Chromosome Counts Database (CCDB) – A community resource of plant chromosome numbers. *The New Phytologist*, 206(1), 19–26. <https://doi.org/10.1111/nph.13191>
- Richardson, S. S. (2013). *Sex itself: The search for male & female in the human genome*. University of Chicago Press.
- Rifkin, J. L., Beaudry, F. E. G., Humphries, Z., ... Wright, S. I. (2020). Widespread recombination suppression facilitates plant sex chromosome evolution. *Molecular Biology and Evolution*, 38(3), 1018–1030. <https://doi.org/10.1093/molbev/msaa271>
- Rifkin, J. L., Hnatovska, S., Yuan, M., ... Wright, S. I. (2022). Recombination landscape dimorphism and sex chromosome evolution in the dioecious plant *Rumex hastatulus*. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 377(1850), 20210226. <https://doi.org/10.1098/rstb.2021.0226>
- Rodrigues, N., Studer, T., Dufresnes, C., & Perrin, N. (2018). Sex-chromosome recombination in common frogs brings water to the fountain-of-youth. *Molecular Biology and Evolution*, 35(4), 942–948. <https://doi.org/10.1093/molbev/msy008>
- Rodríguez-González, R., Gutiérrez, M. L., Fuentes, I., ... Garcia, S. (2023). Release 4.0 of the Plant rDNA database: A database on plant ribosomal DNA loci number, their position, and organization: An information source for comparative cytogenetics. *Methods in Molecular Biology (Clifton, N.J.)*, 2703, 237–245. [https://doi.org/10.1007/978-1-0716-3389-2\\_18](https://doi.org/10.1007/978-1-0716-3389-2_18)
- Román-Palacios, C., Medina, C. A., Zhan, S. H., & Barker, M. S. (2021). Animal chromosome counts reveal a similar range of chromosome numbers but with less polyploidy in animals compared to flowering plants. *Journal of Evolutionary Biology*, 34(8), 1333–1339. <https://doi.org/10.1111/jeb.13884>

- Ross, L., Blackmon, H., Lorite, P., ... Hardy, N. B. (2015). Recombination, chromosome number and eusociality in the Hymenoptera. *Journal of Evolutionary Biology*, 28(1), 105–116. <https://doi.org/10.1111/jeb.12543>
- Ruiz-Herrera, A., & Waters, P. D. (2022). Fragile, unfaithful and persistent Ys—On how meiosis can shape sex chromosome evolution. *Heredity*, 129(1), 22–30. <https://doi.org/10.1038/s41437-022-00532-2>
- Sabath, N., Itescu, Y., Feldman, A., ... Valenzuela, N. (2016). Sex determination, longevity, and the birth and death of reptilian species. *Ecology and Evolution*, 6(15), 5207–5220. <https://doi.org/10.1002/ece3.2277>
- Sacchi, B., Humphries, Z., Kruzlicová, J., ... Wright, S. I. (2024). Phased assembly of Neo-sex chromosomes reveals extensive Y degeneration and rapid genome evolution in *Rumex hastatulus*. *Molecular Biology and Evolution*, 41(4), msae074. <https://doi.org/10.1093/molbev/msae074>
- Sahara, K., Yoshido, A., & Traut, W. (2012). Sex chromosome evolution in moths and butterflies. *Chromosome Research*, 20(1), 83–94. <https://doi.org/10.1007/s10577-011-9262-z>
- Sardell, J. M., & Kirkpatrick, M. (2020). Sex differences in the recombination landscape. *The American Naturalist*, 195(2), 361–379. <https://doi.org/10.1086/704943>
- Sarre, S. D., Georges, A., & Quinn, A. (2004). The ends of a continuum: Genetic and temperature-dependent sex determination in reptiles. *BioEssays*, 26(6), 639–645. <https://doi.org/10.1002/bies.20050>
- Sasson, D. A., & Ryan, J. F. (2017). A reconstruction of sexual modes throughout animal evolution. *BMC Evolutionary Biology*, 17(1), 242. <https://doi.org/10.1186/s12862-017-1071-3>
- Satomura, K., Osada, N., & Endo, T. (2019). Achiasmy and sex chromosome evolution. *Ecological Genetics and Genomics*, 13, 100046. <https://doi.org/10.1016/j.egg.2019.100046>
- Saunders, P. A., & Veyrunes, F. (2021). Unusual mammalian sex determination systems: a cabinet of curiosities. *Genes*, 12(11), 1770. <https://doi.org/10.3390/genes12111770>
- Schärer, L., Rowe, L., & Arnqvist, G. (2012). Anisogamy, chance and the evolution of sex roles. *Trends in Ecology & Evolution*, 27(5), 260–264. <https://doi.org/10.1016/j.tree.2011.12.006>
- Schartl, M., Georges, A., & Marshall Graves, J. A. (2023). Polygenic sex determination in vertebrates – Is there any such thing? *Trends in Genetics: TIG*, 39(4), 242–250. <https://doi.org/10.1016/j.tig.2022.12.002>
- Schlupp, I. (2005). The evolutionary ecology of gynogenesis. *Annual Review of Ecology, Evolution, and Systematics*, 36(1), 399–417. <https://doi.org/10.1146/annurev.ecolsys.36.102003.152629>
- Schmid, M., Nanda, I., Steinlein, C., ... Haaf, T. (1991). Sex-determining mechanisms and sex chromosomes in amphibia. In D. M. Green, & S. K. Sessions (Eds.), *Amphibian cytogenetics and evolution* (pp. 393–430). Academic Press Inc.
- Schneider, M. C., Mattos, V. F., & Cella, D. M. (2024). The scorpion cytogenetic database. [www.arthropodacytogenetics.bio.br/scorpiondatabase](http://www.arthropodacytogenetics.bio.br/scorpiondatabase)
- Schroeder, A. L., Metzger, K. J., Miller, A., & Rhen, T. (2016). A novel candidate gene for temperature-dependent sex determination in the common snapping turtle. *Genetics*, 203(1), 557–571. <https://doi.org/10.1534/genetics.115.182840>
- Schultz, R. J. (1969). Hybridization, unisexuality, and polyploidy in the teleost *Poeciliopsis* (Poeciliidae) and other vertebrates. *The American Naturalist*, 103(934), 605–619. <https://doi.org/10.1086/282629>
- Schwaegerle, K. E. (2005). Quantitative genetic analysis of plant growth: Biases arising from vegetative propagation. *Evolution; International Journal of Organic Evolution*, 59(6), 1259–1267.
- Schwander, T., Henry, L., & Crespi, B. J. (2011). Molecular evidence for ancient asexuality in timema stick insects. *Current Biology: CB*, 21(13), 1129–1134. <https://doi.org/10.1016/j.cub.2011.05.026>
- Schwander, T., & Oldroyd, B. P. (2016). Androgenesis: Where males hijack eggs to clone themselves. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 371(1706), 20150534. <https://doi.org/10.1098/rstb.2015.0534>
- Schwanz, L. E., Georges, A., Holleley, C. E., & Sarre, S. D. (2020). Climate change, sex reversal and lability of sex-determining systems. *Journal of Evolutionary Biology*, 33(3), 270–281. <https://doi.org/10.1111/jeb.13587>
- Sember, A., Nguyen, P., Perez, M. F., ... Cioffi, M. de B. (2021). Multiple sex chromosomes in teleost fishes from a cytogenetic perspective: state of the art and future challenges. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 376(1833), 20200098. <https://doi.org/10.1098/rstb.2020.0098>
- Shanahan, C. M., & Hayman, D. L. (1990). Synaptonemal complex formation in male scorpions exhibiting achiasmate meiosis and structural heterozygosity. *Genome*, 33(6), 914–926. <https://doi.org/10.1139/g90-138>
- Shi, M. -W., Zhang, N. -A., Shi, C. -P., ... Chen, Z. -X. (2019). SAGD: A comprehensive sex-associated gene database from transcripts. *Nucleic Acids Research*, 47(D1), D835–D840. <https://doi.org/10.1093/nar/gky1040>
- Smith, J. J., Timoshevskiy, V. A., & Saraceno, C. (2021). Programmed DNA elimination in vertebrates. *Annual Review of Animal Biosciences*, 9(1), 173–201. <https://doi.org/10.1146/annurev-animal-061220-023220>
- Sochorová, J., Garcia, S., Gálvez, F., ... Kovařík, A. (2018). Evolutionary trends in animal ribosomal DNA loci: Introduction to a new online database. *Chromosoma*, 127(1), 141–150. <https://doi.org/10.1007/s00412-017-0651-8>
- Speijer, D., Lukeš, J., & Eliáš, M. (2015). Sex is a ubiquitous, ancient, and inherent attribute of eukaryotic life. *Proceedings of the National Academy of Sciences of the United States of America*, 112(29), 8827–8834. <https://doi.org/10.1073/pnas.1501725112>
- Štáhlavský, F. (2022). *The pseudoscorpion cytogenetic database*. [www.arthropodacytogenetics.bio.br/pseudoscorpiondatabase](http://www.arthropodacytogenetics.bio.br/pseudoscorpiondatabase)
- Stevens, N. M. (1905). *Studies in spermatogenesis* (Vol. 1). Carnegie Institution of Washington.
- Stevens, N. M. (1906). *Studies in spermatogenesis* (Part 2) (Vol. 2). Carnegie Institution of Washington.
- Stöck, M., Dedukh, D., Reifová, R., ... Janko, K. (2021a). Sex chromosomes in meiotic, hemiclinal, clonal and polyploid hybrid vertebrates: Along the “extended speciation continuum.”. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 376(1833), 20200103. <https://doi.org/10.1098/rstb.2020.0103>
- Stöck, M., Kratochvíl, L., Kuhl, H., ... Guiguen, Y. (2021b). A brief review of vertebrate sex evolution with a pledge for integrative research: Towards “sexomics. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 376(1832), 20200426. <https://doi.org/10.1098/rstb.2020.0426>
- Sylvester, T., Hjelm, C. E., Hanrahan, S. J., ... Blackmon, H. (2020). Lineage-specific patterns of chromosome evolution are the rule not the exception in Polyneoptera insects. *Proceedings Biological Sciences*, 287(1935), 20201388. <https://doi.org/10.1098/rspb.2020.1388>
- Thomas, C. G., Woodruff, G. C., & Haag, E. S. (2012). Causes and consequences of the evolution of reproductive mode in *Caenorhabditis* nematodes. *Trends in Genetics: TIG*, 28(5), 213–220. <https://doi.org/10.1016/j.tig.2012.02.007>
- Thuy, B., Numberger-Thuy, L. D., Härer, J., ... Schweigert, G. (2024). Fossil evidence for the ancient link between clonal fragmentation, six-fold symmetry and an epizoa lifestyle in asterozoan echinoderms. *Proceedings Biological Sciences*, 291(2023), 20232832. <https://doi.org/10.1098/rspb.2023.2832>
- Tilquin, A., & Kokko, H. (2016). What does the geography of parthenogenesis teach us about sex? *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*, 371(1706), 20150538. <https://doi.org/10.1098/rstb.2015.0538>
- Titley, M. A., Snaddon, J. L., & Turner, E. C. (2017). Scientific research on animal biodiversity is systematically biased towards vertebrates and temperate regions. *PLoS One*, 12(12), e0189577. <https://doi.org/10.1371/journal.pone.0189577>

- Toups, M. A., & Vicoso, B. (2023). The X chromosome of insects likely predates the origin of class Insecta. *Evolution; International Journal of Organic Evolution*, 77(11), 2504–2511. <https://doi.org/10.1093/evolut/qpaa169>
- Tree of Sex Consortium (2014). Tree of Sex: A database of sexual systems. *Scientific Data*, 1, 140015.
- Tsurusaki, N., Svojanovská, H., Schöenhofer, A., & Šťáhlavský, F. (2022). *The harvestmen cytogenetic database*. <http://arthropodacytogenetics.bio.br/harvestmendatabase>
- Turelli, M., & Orr, H. A. (1995). The dominance theory of Haldane's rule. *Genetics*, 140(1), 389–402. <https://doi.org/10.1093/genetics/140.1.389>
- Turner, L. M., & Harr, B. (2014). Genome-wide mapping in a house mouse hybrid zone reveals hybrid sterility loci and Dobzhansky-Muller interactions. *eLife*, 3, 3. <https://doi.org/10.7554/eLife.02504>
- Valenzuela, N., & Lance, V. (2004). *Temperature dependent sex determination in vertebrates*. Smithsonian Books.
- Valenzuela, N., Litterman, R., Neuwald, J. L., ... Litzgus, J. D. (2019). Extreme thermal fluctuations from climate change unexpectedly accelerate demographic collapse of vertebrates with temperature-dependent sex determination. *Scientific Reports*, 9(1), 1–11.
- Van Dijk, P. (2009). Apomixis: Basics for non-botanists. In Schön, D. I., Martens, K. (Eds.), *Lost sex* (pp. 47–62). Springer Netherlands.
- Vara, C., Paytuví-Gallart, A., Cuartero, Y., ... Ruiz-Herrera, A. (2019). Three-dimensional genomic structure and cohesin occupancy correlate with transcriptional activity during spermatogenesis. *Cell Reports*, 28(2), 352–367.e9. <https://doi.org/10.1016/j.celrep.2019.06.037>
- Vara, C., Paytuví-Gallart, A., Cuartero, Y., ... Ruiz-Herrera, A. (2021). The impact of chromosomal fusions on 3D genome folding and recombination in the germ line. *Nature Communications*, 12(1), 2981. <https://doi.org/10.1038/s41467-021-23270-1>
- Veltsos, P., Shinde, S., & Ma, W.-J. (2024). Sex chromosome evolution: The classical paradigm and so much beyond. *arXiv*, <http://arxiv.org/abs/2408.12034>, August 2024, preprint: not peer reviewed.
- Vicoso, B. (2019). Molecular and evolutionary dynamics of animal sex-chromosome turnover. *Nature Ecology & Evolution*, 3(12), 1632–1641. <https://doi.org/10.1038/s41559-019-1050-8>
- Vicoso, B., & Bachtrog, D. (2015). Numerous transitions of sex chromosomes in Diptera. *PLoS Biology*, 13(4), e1002078. <https://doi.org/10.1371/journal.pbio.1002078>
- Villarreal, J. C., & Renner, S. S. (2013). Correlates of monoicy and dioicy in hornworts, the apparent sister group to vascular plants. *BMC Evolutionary Biology*, 13(1), 239. <https://doi.org/10.1186/1471-2148-13-239>
- Wang, J., & Davis, R. E. (2014). Programmed DNA elimination in multicellular organisms. *Current Opinion in Genetics & Development*, 27, 26–34. <https://doi.org/10.1016/j.gde.2014.03.012>
- Wang, J., Tao, W., Kocher, T. D., & Wang, D. (2024). Sex chromosome turnover and biodiversity in fishes. *Journal of genetics and genomics = Yi chuan xue bao*, 51(12), 1351–1360. <https://doi.org/10.1016/j.jgg.2024.08.008>
- Weber, C., Zhou, Y., Lee, J. G., ... Capel, B. (2020). Temperature-dependent sex determination is mediated by pSTAT3 repression of Kdm6b. *Science*, 368(6488), 303–306. <https://doi.org/10.1126/science.aaz4165>
- Weeks, S. C. (2012). The role of androdioecy and gynodioecy in mediating evolutionary transitions between dioecy and hermaphroditism in the animalia: Reproductive transitions in the animalia. *Evolution; International Journal of Organic Evolution*, 66(12), 3670–3686. <https://doi.org/10.1111/j.1558-5646.2012.01714.x>
- Weeks, S. C., Benvenuto, C., & Reed, S. K. (2006). When males and hermaphrodites coexist: A review of androdioecy in animals. *Integrative and Comparative Biology*, 46(4), 449–464. <https://doi.org/10.1093/icb/icj048>
- Weir, W., Capewell, P., Foth, B., ... MacLeod, A. (2016). Population genomics reveals the origin and asexual evolution of human infective trypanosomes. *eLife*, 5, e11473. <https://doi.org/10.7554/eLife.11473>
- Werren, J. H., Baldo, L., & Clark, M. E. (2008). *Wolbachia*: Master manipulators of invertebrate biology. *Nature Reviews. Microbiology*, 6(10), 741–751. <https://doi.org/10.1038/nrmicro1969>
- Westneat, D. F., & Craig Sargent, R. (1996). Sex and parenting: The effects of sexual conflict and parentage on parental strategies. *Trends in Ecology & Evolution*, 11(2), 87–91. [https://doi.org/10.1016/0169-5347\(96\)81049-4](https://doi.org/10.1016/0169-5347(96)81049-4)
- White, M. J. D. (1945). *Animal cytology and evolution* (3rd ed.). Cambridge University Press.
- Whiteley, S. L., Georges, A., Weisbecker, V., ... Holleley, C. E. (2021). Ovotestes suggest cryptic genetic influence in a reptile model for temperature-dependent sex determination. *Proceedings Biological Sciences*, 288(1943), 20202819. <https://doi.org/10.1098/rspb.2020.2819>
- Whiteley, S. L., Wagner, S., Holleley, C. E., ... Georges, A. (2022). Truncated *jarid2* and *kdm6b* transcripts are associated with temperature-induced sex reversal during development in a dragon lizard. *Science Advances*, 8(16), eabk0275. <https://doi.org/10.1126/sciadv.abk0275>
- Whiteley, S. L., Weisbecker, V., Georges, A., ... Holleley, C. E. (2018). Developmental asynchrony and antagonism of sex determination pathways in a lizard with temperature-induced sex reversal. *Scientific Reports*, 8(1), 14892. <https://doi.org/10.1038/s41598-018-33170-y>
- Wiggins, J. M., Santoyo-Brito, E., Scales, J. B., & Fox, S. F. (2020). Gene dose indicates presence of sex chromosomes in collared lizards (*Crotaphytus collaris*), a species with temperature-influenced sex determination. *Herpetologica*, 76(1), 27–30. <https://doi.org/10.1655/herpetologica-d-19-00036>
- Wright, C. J., Stevens, L., Mackintosh, A., ... Blaxter, M. (2024). Comparative genomics reveals the dynamics of chromosome evolution in Lepidoptera. *Nature Ecology & Evolution*, 8(4), 777–790. <https://doi.org/10.1038/s41559-024-02329-4>
- Wright, S. I., Kalisz, S., & Slotte, T. (2013). Evolutionary consequences of self-fertilization in plants. *Proceedings Biological Sciences*, 280(1760), 20130133. <https://doi.org/10.1098/rspb.2013.0133>
- Ye, Z., Bishop, T., Wang, Y., ... Lynch, M. (2023). Evolution of sex determination in crustaceans. *Marine Life Science & Technology*, 5(1), 1–11. <https://doi.org/10.1007/s42995-023-00163-4>
- Yusa, Y. (2007). Causes of variation in sex ratio and modes of sex determination in the Mollusca—An overview. *American Malacological Bulletin*, 23(1), 89–98. <https://doi.org/10.4003/0740-2783-23.1.89>
- Zaccanti, F., & Farne', P. (1986). Observations on fissiparity and spontaneous regeneration in two strains of agamic planarians (*Dugesia gonocephala* and *Dugesia tigrina*). *The Journal of Experimental Zoology*, 238(3), 319–324. <https://doi.org/10.1002/jez.1402380306>
- Zaidem, M. L., Groen, S. C., & Purugganan, M. D. (2019). Evolutionary and ecological functional genomics, from lab to the wild. *The Plant Journal: for Cell and Molecular Biology*, 97(1), 40–55. <https://doi.org/10.1111/tpj.14167>
- Zayed, A., & Packer, L. (2005). Complementary sex determination substantially increases extinction proneness of haplodiploid populations. *Proceedings of the National Academy of Sciences of the United States of America*, 102(30), 10742–10746. <https://doi.org/10.1073/pnas.0502271102>